



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, C12P 19/34, G01N 33/54, C07H 21/02, 21/04, C07K 15/26, 15/28		A1	(11) International Publication Number: WO 99/11824
			(43) International Publication Date: 11 March 1999 (11.03.99)
(21) International Application Number: PCT/US98/18535		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 8 September 1998 (08.09.98)		Published <i>With international search report.</i>	
(30) Priority Data: 60/057,655 5 September 1997 (05.09.97) US 60/093,297 17 July 1998 (17.07.98) US 09/148,503 4 September 1998 (04.09.98) US			
(71)(72) Applicants and Inventors: TANZI, Rudolph, E. [US/US]; 3 Oceanside Drive, Hull, MA 02045 (US). HYMAN, Bradley, T. [US/US]; 23 Bayview Avenue, Swampscott, MA 01907 (US). REBECK, George, W. [US/US]; Apartment 1, 93 Concord Avenue, Somerville, MA 02143 (US).			
(74) Agents: GOLDSTEIN, Jorge, A. et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US).			
(54) Title: AN ALPHA-2 MACROGLOBULIN DIAGNOSTIC TEST			
(57) Abstract			
<p>The disclosed invention relates to a diagnostic method of Alzheimer's disease based on genotyping the Alpha-2-Macroglobulin locus. A statistically significant correlation was found between inheritance of particular alleles of the Alpha-2-Macroglobulin gene and the occurrence of Alzheimer's disease. The diagnostic method involves the isolation of nucleic acid from an individual and subsequent genotyping by means such as sequencing or restriction fragment length polymorphism analysis. The invention also provides a means of genotype analysis through protein isotyping Alpha-2-Macroglobulin variant proteins. Finally, kits for nucleic acid analysis or protein analysis are taught.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

An Alpha-2 Macroglobulin Diagnostic Test

Background of the Invention

5 ***Statement as to Rights to Inventions Made Under Federally-Sponsored Research and Development***

At least part of the work performed during development of this invention utilized Grant
Nos. P50AG05134 and R01AG12406, from the National Institute of Aging, and Grant No.
10 5U01 MH51066, from the National Institute of Mental Health. The U.S. Government has certain
rights in this invention.

Field of the Invention

15 This invention relates to the area of medical genetics. Specifically, the invention provides
a genetic test to identify carriers of specific alleles of the Alpha-2-Macroglobulin gene, the
occurrence of which are genetically linked, with high statistical significance, to the condition
known as Alzheimer's disease.

20 ***Related Art***

Alpha-2-Macroglobulin (α_2 M) is a major serum pan-protease inhibitor which inhibits all
four classes of proteases by a unique steric trapping mechanism (Borth, W., *FASEB* 6:3345-3353
(1992)). It is the major representative of a family of plasma proteins, most of which possess a
25 unique internal cyclic thiol ester bond, referred to as the α -macroglobulins (Borth, W., *FASEB*
6:3345-3353 (1992)). Although it functions against a wide variety of human and bacterial
endopeptidases (Starkey, P. M. and Barret, A. J., *Proteinases in mammalian cells and tissues*,
Barret, A. J. (ed.), North-Holland, New York (pub), pp. 663-696, (1977)), a possible role in
pathogenesis is not yet known. *In vitro* studies have provided a great deal of information
30 regarding protein structure and inhibitory functions (Scottrup-Jensen, L., *et al.*, *J. Biol. Chem.*
259:8318-8327 (1984); Feldman, S. R., *et al.*, *Proc. Natl. Acad. Sci. USA* 82:5700-5704, 1985;
Kan, C. C., *et al.*, *Proc. Natl. Acad. Sci. USA* 82:2282-2286 (1985); Bretaudiere, J. P., *et al.*,

5 *Proc. Natl. Acad. Sci. USA* 85:1437-1441 (1988); Van leuven, F., *et al.*, *J. Biol. Chem.* 261:11369-11373 (1986)), and at the molecular level, it is known to be part of a related gene cluster on human chromosome 12p (Devriendt, K., *et al.*, *Gene* 81:325-334, 1989) that also includes the α_2 M pseudogene and the pregnancy-zone gene. Recent evidence suggests that α_2 M may be associated with certain morphological characteristics of Alzheimer's Disease (AD).

10 AD is a neurodegenerative disorder characterized by a global decline in mental function, memory and acquired intellectual skills. It is the most common form of dementia occurring in mid-to late-life and is a major cause of disability and death in the elderly (Price, L. and Sisodia, S., *Annu. Rev. Neurosci.* 21:479-505 (1998)). The appearance of AD in the population according to age is variable, often being categorized into early and late onset forms. In the general
15 population, 40% of the brains of normal individuals show some evidence of A β deposits, indicating a subclinical prevalence, and in the population 60 years and older, AD is diagnosed in 10% of the population.

AD is a genetically heterogenous disorder. Early-onset familial AD (EO-FAD) is inherited as an autosomal dominant disorder involving defects in at least three genes, presenilin I (*PSEN1*)
20 on chromosome 14 (Sherrington, R., *et al.*, *Nature* 375:754-760 (1995)), presenilin 2 (*PSEN2*) on chromosome 1 (Levy-Lehad, E., *et al.*, *Science* 269:973-977 (1995); Rogaev, E. I., *et al.*, *Nature* 376:775-778 (1995)), and β -amyloid precursor protein (*APP*) on chromosome 21 (Tanzi, R. E., *et al.*, *Science* 235:880-884 (1987); Goate, A. M., *et al.*, *Nature* 349:704-706 (1991)). These genes account for roughly 30-40% of EO-FAD (Hardy J., *Trends Neurosci.* 20:154-159
25 (1997); Cruts, M., *et al.*, *Hum. Mol. Genet.* 7:43-51 (1998); Blacker, D. and Tanzi, R. E., *Arch Neurol* 55:294-296 (1998)).

Late-onset AD (LOAD) has been associated with the *APOE- ϵ 4* allele of apolipoprotein E (*APOE*) located on chromosome 19 (Strittmatter, W. J., *et al.*, *Proc. Natl. Acad. Sci. (USA)* 90:1977-1981 (1993); Saunders, A. M., *et al.*, *Neurology* 43:1467-1472 (1993)). Inheritance of
30 *APOE- ϵ 4* lowers the age of onset of AD in a dose-dependent manner and is associated with cases of AD occurring from 40 to 90 years. However it has the greatest impact as a risk factor for onset in the 60's (Blacker, D., *et al.*, *Neurology* 48:139-147 (1997); Farrer, L. A., *et al.*, *JAMA* 278:1349-1356 (1997)). More recently, the gene encoding the major neuronal receptor for apoE, the low density lipoprotein receptor-related protein (LRP), located on the long arm of

5 chromosome 12, was shown to be associated with LOAD (Kang, D. E., *et al.*, *Neurology* 49:56-61 (1997)).

Neuropathological hallmarks of AD include abundant neurofibrillary tangles (NFT), and β -amyloid deposition in cerebral blood vessels and in senile plaques (SP). Deposited amyloid is composed principally of the 40-42 residue β -amyloid protein ($A\beta$) (Glennner *et al.*, *Biochem, Biophys Res. Commun.* 120:885-890 (1984)), which is a proteolytic fragment of the β -amyloid precursor protein (APP) (Kang *et al.*, *Nature* 325:733-736 (1987)). Pathogenesis is thought to develop in AD patients when the $A\beta$ protein is organized into neurotoxic fibrils. *In vitro* studies demonstrate that $A\beta$ is generated as a soluble peptide during cellular metabolism (Haas, C., *et al.*, *Nature* 359:322-325, 1992; Shoji, M., *et al.*, *Science* 258:126-129 (1992)). Several additional plaque-associated proteins are known to promote the *in vitro* formation of $A\beta$ fibrils, suggesting the possibility that these proteins regulate $A\beta$ aggregation *in vivo* (Ma, J., *et al.*, *Nature* 372: 92-94 (1994); Eriksson, S., *et al.*, *Proc Natl. Acad. Sci. USA* 92:2313-2317 (1995)). Because $A\beta$ deposition likely plays a role in the development of AD, a great deal of research is focused on the synthesis, degradation and clearance of this protein from the affected tissues.

Furthermore, given the potential importance of the APP and $A\beta$ proteins in AD pathogenesis, research is also focused on providing a genetic link between mutations in the APP gene and the occurrence of Alzheimer's disease. Mutations in APP close to or in the $A\beta$ domain are known (Goate *et al.*, *Nature* 349:704-706 (1991); Levy *et al.*, *Science* 248:1124-1126 (1990); Murrell *et al.*, *Science* 254:97-99 (1991); Hendricks *et al.*, *Nature Genet.* 1:218-221 (1992); Chartier-Harlin, M. *et al.*, *Nature* 353:844 (1991); Mullan, M., *et al.*, *Nature Genet.* 1:345 (1992)). In some instances, linkage of the mutation with the occurrence of familial Alzheimer's disease (FAD) was possible.

Also implicated in the development of AD is the possibility that there exists an imbalance between proteases and protease inhibitors which affects normal amyloid metabolism. Antibodies to ubiquitin strongly react with the characteristic amyloid neurofibrillar tangles (Tabaton, M., *et al.*, *Proc. Natl. Acad. Sci.* 88:2098-2102 (1991)). The $A\beta$ peptide is generated proteolytically from APP, and proteases cathepsin B and D and protease inhibitors such as protease nexin 1 and α 1-antichymotrypsin were found in amyloid plaques (Cataldo, A. M. and Nixon, R. A., *Proc.*

- 5 *Natl. Acad. Sci.* 87:3861-3865, 1991; Rosenblatt, D. E., *et al.*, *Ann. Neurol.* 26:628-634 (1989); Abraham, C. R., *et al.*, *Cell* 52:487-501 (1988)).

Alpha-2-macroglobulin is another protease inhibitor implicated in neural tissue metabolism. Several lines of evidence link the α_2 M protein to morphological properties associated with AD patient cerebra. One study in support of a role in neural metabolism identified α_2 M as an astroglia-derived neurite-promoting factor for cultured neurons from the rat central nervous system (Mori, T., *et al.*, *Brain Res.* 527:55-61 (1990)). In addition, α_2 M has been shown to attenuate the fibrillogenesis and neurotoxicity of A β (Hughes, S. R., *et al.*, *Proc. Natl. Acad. Sci. (USA)* 95:3275-3280 (1998); Du, Y., *et al.*, *J. Neurochem.* 70:1182-1188 (1998); Zhang, Z., *et al.*, *Amyloid: Int. J. Exp. Clin. Invest.* 3:156-161 (1996)). Also suggestive of a connection between α_2 M and AD is that α_2 M tightly binds the A β peptide (Du, Y., *et al.*, *J. Neurochem.* 69:299-305 (1997); Hughes, S. R., *et al.*, *Proc. Natl. Acad. Sci. (USA)* 95:3275-3280 (1998)), and has been shown to mediate the clearance of A β via endocytosis through LRP (Narita, M., *et al.*, *J. Neurochem.* 69: 1904-1911 (1997)). Furthermore, two independent studies found that an α_2 M/protease complex is capable of degrading the A β peptide (Zhang, Z., *et al.*, *Amyloid: Int. J. Exp. Clin. Invest.* 3:156-161 (1996); Qiu, W. Q., *et al.*, *J. Biol. Chem.* 271(14):8443-8451 (1996)). In addition, α_2 M has been localized to senile plaques (SP) in AD (Bauer, J., *et al.*, *FEBS* 285:111-114 (1991); Van Gool, D., *et al.*, *Neurobiology of Aging* 14:233-237 (1993); Rebeck, G. W., *et al.*, *Ann. Neurol.* 37:211-217 (1995)). Van Gool *et al.*, suggest that expression of α_2 M could be associated with neurofibrillary changes in senile plaques in AD. (Van Gool, D., *et al.*, *Neurobiology of Aging* 14:233-237 (1993)). An immunocytochemistry study of cerebra of AD patients using two different monoclonal antibodies specific for α_2 M found localization in neuritic-type plaques but not preamyloid or burned-out type plaques (Van Gool, D., *et al.*, *Neurobiology of Aging* 14:233-237 (1993)). Because immunoreactivity was associated with microglia in the outer border of the neuritic plaque, Van Gool *et al.* suggest that α_2 M could be a marker for an inflammatory process in the plaques (Van Gool, D., *et al.*, *Neurobiology of Aging* 14:233-237 (1993)).

The human α_2 M gene has been cloned and several mutants have been identified. (Poller, W., *et al.*, *Human Genetics* 88:313-319 (1992); Matthijs, G., and Marynen, P., *Nucl. Acids Res.* 19(18):5102 (1991)). One mutation represented a bi-allelic deletion polymorphism in which 5

5 bases were deleted at positions -7 to -3 of the 5' splice acceptor site of exon 18 (Matthijs, G., and Marynen, P., *Nucl. Acids Res.* 19(18):5102 (1991)) (referred to as the *A2M-2* allele). The *A2M-2* allele, and the wild type *A2M* allele (referred to as the *A2M-1* allele) were found at allele frequencies of 0.18 and 0.82 respectively (Matthijs, G., and Marynen, P., *Nucl. Acids Res.* 19(18):5102 (1991)). The biological consequences of this deletion mutation have not yet been
10 reported, but it is known that exon 18 encodes "exon II" of the bait domain of α_2M , which is used to trap proteases.

Another mutation represented a simple polymorphism 25 amino acids downstream from the thiolester site of the protein that interchanges Val¹⁰⁰⁰ (GTC) and Ile¹⁰⁰⁰ (ATC) (Poller, W., *et al.*, *Human Genetics* 88:313-319 (1992) (numbering is based on the cDNA sequence which
15 includes a 24 amino acid signal peptide; this amino acid corresponds to Val/Ile⁹⁷⁶ in the mature protein). The Val, or G allele, and the Ile, or A allele, were found at allele frequencies of 0.30 and 0.70, respectively (Poller, W., *et al.*, *Human Genetics* 88:313-319 (1992)). No difference in α_2M serum levels was associated with the two alleles. This polymorphism is interesting because the mutation occurs near the thiolester active site of the molecule.

20 Family, twin, and population data all suggest that genes involved in AD remain to be identified (Lautenschlager, N. T., *et al.*, *Neurology* 46(3):641-50 (1996); Bergem, A., *et al.*, *Arch. Gen. Psychiatry* 54:264-270 (1997); Payami, H., *et al.*, *Am J Hum Genet* 60:948-956 (1997)) and several candidates have been reported (for a review, see Hardy J., *Trends Neurosci.* 20:154-159 (1997); Cruts, M., *et al.*, *Hum. Mol. Genet.* 7:43-51 (1998); and Blacker, D., and
25 Tanzi, R.E., *Arch. Neurol.* 55:294-296 (1998)). The identification of specific genes or alleles thereof linked to the development of AD will provide a greater understanding of the mechanisms behind the disease process, a screening method for detection of at-risk individuals, and may lead to effective treatment of this disease. A great deal towards this end has been accomplished already in the study of the *APP* gene. This invention, linking *A2M*, and particularly inheritance
30 of the *A2M-2* and *A2M-G* alleles, to the occurrence of Alzheimer's disease, provides another important resource.

Summary of the Invention

This invention relates to the discovery that particular alleles of *A2M* are linked to the occurrence of Alzheimer's disease. More specifically, it was found that individuals carrying at least one copy of the *A2M-2* allele or who are homozygous for the *A2M-G* allele are disproportionately represented in a population of AD patients as compared to those unaffected by AD.

Based on this discovery, a diagnostic method was developed that characterizes the *A2M* genotype of individuals in the population to assess risk for developing AD. In one embodiment of the invention, *A2M* genotype is determined by isolating nucleic acid from an individual, and analyzing it for the particular *A2M* alleles of interest. Various embodiments of the invention enable analysis from isolated nucleic acid that is DNA or RNA. In a preferred embodiment, restriction fragment length polymorphism (RFLP) analysis is used to determine *A2M* genotype.

In a more preferred embodiment of the invention, the nucleic acid is utilized as a template for the amplification of an *A2M* gene fragment prior to the genotyping analysis. A preferred method is to amplify a fragment that contains the site of the pentanucleotide deletion mutation found in the *A2M-2* allele. Analysis of the product of amplification is preferably accomplished by determining the polynucleotide sequence of the amplified fragment. Other methods of analyzing the amplified fragment include size determination, single-strand conformation polymorphism (SSCP) analysis, and RFLP analysis. Another preferred method is to amplify a fragment that encodes amino acid 1000, the residue affected by the mutation found in the *A2M-G* allele. Analysis of the product of amplification is preferably accomplished by RFLP analysis. Other methods of analyzing the amplified fragment include sequencing or sizing the fragment, and SSCP analysis.

Another embodiment of the invention relates to the use of protein isotyping to assess *A2M* genotype. In one embodiment, an antibody specific for an α_2 M variant lacking exon 18 (α_2 M-2) is used to assess *A2M* genotype through protein-isotyping. In a preferred method of protein isotyping, a second antibody, which is specific for an α_2 M protein having exon 18 (the wild type α_2 M referred to as α_2 M-1), is used as a control. Yet another embodiment of the invention relates to the use of an antibody specific for the Val¹⁰⁰⁰ variant to assess the *A2M-G* genotype through

5 protein isotyping. In a preferred method of protein isotyping, a second antibody, which is specific for the Ile¹⁰⁰⁰ protein, is used as a control. For the protein isotyping methods of the invention, preferably, protein is isolated from an individual and screened with the specific antibodies. More specifically, a more preferred embodiment would utilize Western blot or ELISA technology. In yet another embodiment of the invention an α_2 M electrophoretic mobility assay is used to detect
10 the α_2 M-2 variant to assess *A2M* genotype through protein isotyping.

The invention also relates to diagnostic kits for *A2M* genotyping. Alternative embodiments of the genotyping kit enable screening of nucleic acid or protein, providing a fast and efficient means for Alzheimer's disease risk assessment.

15 *Brief Description of the Figures*

FIG. 1 Map of Chromosome 12 Markers: Figure 1 depicts a map of several chromosome 12 markers. Distances between markers are given in cM. * represents markers used in the traditional linkage analysis techniques described in Example 1 below; @ represents markers
20 used in the analysis described by Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)); \$ represents markers used in both analyses. The letter T inside of a circle represents a telomere of Chromosome 12. The letter C inside of a circle represents the centromere of Chromosome 12.

FIGS. 2A-2D Results of Multipoint Non Parametric Linkage Analysis of
25 **Chromosome 12 Markers to Alzheimer's Disease:** Figures 2A-2D depict the results of a multipoint non-parametric linkage analysis using the program GENEHUNTER (Kruglyak, L., *et al.*, *Am. J. Hum. Genet.* 58:1347-1363 (1996)(software available from Kruglyak *et al.* by anonymous ftp at ftp-genome.wi.mit.edu, or from their World Wide Web site <http://www-genome.wi.mit.edu/ftp/distribution/software/genehunter>)). This was one of several genetic
30 linkage analytic techniques used to determine if there was evidence for linkage in the *A2M* region, and particularly, whether *A2M* might be related to recent reports of linkage to the centromeric region of chromosome 12 (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)). The results were negative, although there was a weak but non-significant signal in Tier 1. For this analysis, the families from the sample described in Example 1 were divided into "tiers" according

5 to the criteria of Pericak-Vance *et al.*: Tier 1 (Figure 2B) is comprised of families in which all affecteds were *APOE-ε4/ε4* (30 families); Tier 2 (Figure 2C) is comprised of families not in Tier 1 in which all affecteds were *APOE-ε4* carriers (131 families); Tier 3 (Figure 2D) is comprised of families in which at least one affected did not carry an *APOE-ε4* allele (126 families). The analysis was also run on the total sample (Figure 2A).

10 **FIG. 3 Alpha-2-Macroglobulin Val¹⁰⁰⁰ Polymorphism Genotyping:** Figure 3 is a photograph of a 2% agarose gel showing *A2M* polymorphism genotypes after restriction digestion with *MboI*. The gel shows a restriction digest pattern with bands representing the G (532 bp fragment) and the A (429 bp fragment) alleles. Genomic DNA isolated from brain tissue and blood was amplified by polymerase chain reaction (PCR) in the presence of oligonucleotide sense
15 primer C23 (SEQ ID NO: 3) and antisense primer AS24 (SEQ ID NO: 4), 10 mM Tris-HCl, 50 mM KCl (pH 8.3), 1.5 mM MgCl₂, 5 mM dNTPs, 5 pmol each primer, and 1.25 U *Taq* DNA polymerase. The PCR was carried out in a touchdown procedure that stepped down the annealing temperature to increase primer specificity as follows: 1 cycle at 94°C-5 min; 4 cycles at 94°C-30 sec, 65°C-30 sec, 72°C-1 min; 4 cycles at 94°C-30 sec, 62°C-30 sec, 72°C-1 min; 4 cycles at
20 94°C-30 sec, 59°C-30 sec, 72°C-1 min; 20 cycles at 94°C-30 sec, 56°C-30 sec, 72°C-1 min; and 1 cycle at 72°C-5 min. For each reaction mixture, 10 units of *MboI* were added to the amplified product of 615 bps and digestion carried out at 37°C for 3 hours, producing fragments of 532 and 429 bps. The digested product was loaded onto a 2% agarose gel treated with ethidium bromide (0.005%) and electrophoresed for 2 hours under constant voltage (150V), which is sufficient to
25 separate the digested product so that the 532 and 429 bp bands can be distinguished. After electrophoresis, DNA fragments were visualized by UV illumination using a Biorad Geldoc system. Incomplete digestion was monitored by looking for the presence of the 615 bp fragment.

Detailed Description of the Preferred Embodiments

Definitions

30 In the description that follows, a number of terms used in recombinant DNA technology are extensively utilized. In order to provide a clearer and consistent understanding of the

5 specification and claims, including the scope to be given such terms, the following definitions are provided.

Polymorphic: As is understood by one of ordinary skill in the art, a nucleic acid molecule is said to be "polymorphic" if it may exist in more than one form. For example, a nucleic acid molecule is said to be polymorphic if it may have more than one specific nucleotide sequence
10 (such as degenerate nucleic acid molecules or genes that may each encode the same protein). More commonly, a nucleic acid molecule is said to be polymorphic if it displays size differences (*i.e.*, differences in length), particularly when comparisons of nucleic acid molecules from different individuals are made. Of course, other definitions of the term "polymorphic" will be apparent to one of ordinary skill and
15 are also encompassed within this definition.

Primer: As used herein "primer" refers to a single-stranded oligonucleotide that is extended by covalent bonding of nucleotide monomers during amplification or polymerization of a DNA molecule. Minisatellite primers used for the amplification of minisatellite dimer, trimer, tetramer, etc., sequences are well-known in the art.

20 **Restriction Fragment Length Polymorphism (RFLP):** As used herein, a restriction fragment polymorphism refers to a polymorphism in the length of restriction fragments produced by digestion of DNA by a particular restriction endonuclease. The genomic DNA of two individuals in a population will differ in sequence at many sites. When these differences occur in the recognition site for a restriction endonuclease, then a polymorphism in the length of restriction
25 fragments produced by digestion of the DNA of the two individuals will result.

Single-Strand Conformation Polymorphism (SSCP): As used herein, a single-strand conformation polymorphism refers to a structural variation in single-stranded DNA due to sequence-dependent intrastrand secondary structure. Secondary structure is determined by the balance between destabilizing thermal forces and weak stabilizing forces. These forces are in turn
30 dependent on the primary structure of the DNA strand, allowing a sequence variation as small as a single base point mutation to lead to structural variations between otherwise identical sequences.

The differences in secondary structure among variants allow the detection of SSCPs by electrophoresis using non-denaturing gels. Under the appropriate conditions, the electrophoretic mobility of the DNA is dependent not only on its length and molecular weight, but also on its

5 conformation. Thus, not only will complementary strands migrate as separate bands on a gel, but small differences in sequence will also alter the mobility of these strands.

Template: The term "template" as used herein refers to a double-stranded or single-stranded nucleic acid molecule which is to be amplified, synthesized or sequenced. In the case of a double-stranded DNA molecule, denaturation of its strands to form a first and a second strand is performed before these molecules may be amplified, synthesized or sequenced. A primer, complementary to a portion of a DNA template is hybridized under appropriate conditions and the DNA polymerase of the invention may then synthesize a DNA molecule complementary to said template or a portion thereof. The newly synthesized DNA molecule, according to the invention, may be equal or shorter in length than the original DNA template. Mismatch incorporation or strand slippage during the synthesis or extension of the newly synthesized DNA molecule may result in one or a number of mismatched base pairs. Thus, the synthesized DNA molecule need not be exactly complementary to the DNA template.

Incorporating: The term "incorporating" as used herein means becoming a part of a DNA molecule or primer.

20 **Amplification:** As used herein "amplification" refers to any *in vitro* method for increasing the number of copies of a nucleotide sequence with the use of a DNA polymerase. Nucleic acid amplification results in the incorporation of nucleotides into a DNA or molecule or primer thereby forming a new DNA molecule complementary to a DNA template. The formed DNA molecule and its template can be used as templates to synthesize additional DNA molecules. As used herein, one amplification reaction may consist of many rounds of DNA replication. DNA amplification reactions include, for example, polymerase chain reactions (PCR). One PCR reaction may consist of 5 to 100 "cycles" of denaturation and synthesis of a DNA molecule.

25 **Oligonucleotide:** "Oligonucleotide" refers to a synthetic or natural molecule comprising a covalently linked sequence of nucleotides which are joined by a phosphodiester bond between the 3' position of the pentose of one nucleotide and the 5' position of the pentose of the adjacent nucleotide.

30 **Nucleotide:** As used herein "nucleotide" refers to a base-sugar-phosphate combination. Nucleotides are monomeric units of a nucleic acid sequence (DNA and RNA). The term nucleotide includes deoxyribonucleoside triphosphates such as dATP, dCTP, dITP, dUTP, dGTP,

5 dTTP, or derivatives thereof. Such derivatives include, for example, [α S]dATP, 7-deaza-dGTP and 7-deaza-dATP. The term nucleotide as used herein also refers to dideoxyribonucleoside triphosphates (ddNTPs) and their derivatives. Illustrated examples of dideoxyribonucleoside triphosphates include, but are not limited to, ddATP, ddCTP, ddGTP, ddITP, and ddTTP. According to the present invention, a "nucleotide" may be unlabeled or detectably labeled by well
10 known techniques. Detectable labels include, for example, radioactive isotopes, fluorescent labels, chemiluminescent labels, bioluminescent labels and enzyme labels.

Thermostable: As used herein "thermostable" refers to a DNA polymerase which is resistant to inactivation by heat. DNA polymerases synthesize the formation of a DNA molecule complementary to a single-stranded DNA template by extending a primer in the 5'-to-3' direction.
15 This activity for mesophilic DNA polymerases may be inactivated by heat treatment. For example, T5 DNA polymerase activity is totally inactivated by exposing the enzyme to a temperature of 90°C for 30 seconds. As used herein, a thermostable DNA polymerase activity is more resistant to heat inactivation than a mesophilic DNA polymerase. However, a thermostable DNA polymerase does not mean to refer to an enzyme which is totally resistant to
20 heat inactivation and thus heat treatment may reduce the DNA polymerase activity to some extent. A thermostable DNA polymerase typically will also have a higher optimum temperature than mesophilic DNA polymerases.

Hybridization: The terms "hybridization" and "hybridizing" refer to the pairing of two complementary single-stranded nucleic acid molecules (RNA and/or DNA) to give a double-
25 stranded molecule. As used herein, two nucleic acid molecules may be hybridized, although the base pairing is not completely complementary. Accordingly, mismatched bases do not prevent hybridization of two nucleic acid molecules provided that appropriate conditions, well known in the art, are used. In the present invention, the term "hybridization" refers particularly to hybridization of an oligonucleotide to a DNA template molecule.

30 Other terms used in the fields of recombinant DNA technology and molecular and cell biology as used herein will be generally understood by one of ordinary skill in the applicable arts.

This invention relates to the finding that particular genotypes of the *A2M* locus may be linked to the occurrence of Alzheimer's disease. More specifically, it was found that individuals

5 carrying at least one copy of the *A2M-2* allele or who are homozygous for the *A2M-G* allele are more highly represented in AD patients than in the population as a whole.

Therefore, a first embodiment of the invention is to utilize *A2M* genotyping as a diagnostic method for determining risk-assessment for the onset of AD. To that end, cells, tissues, organs or other samples from an individual are obtained in order to isolate nucleic acid or protein.

10 Preparation of the nucleic acid is done by methods that are well-known in the art (See, e.g., Maniatis, T., *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press (1989); Kaufman, P.B., *et al.*, *Handbook of Molecular and Cellular Methods in Biology and Medicine*, Boca Raton, Florida: CRC Press (1995)). In one embodiment of the invention, the nucleic acid isolated is DNA. In another embodiment the

15 nucleic acid is RNA, which is subsequently converted to complimentary DNA (cDNA) prior to processing and analysis by means of reverse transcription using procedures well known in the art (see, e.g., Unit 5.5 in Ausubel, F.M. *et al.*, eds., *Current Protocols in Molecular Biology*: John Wiley and Sons, Inc. (1997)). The nucleic acid samples thus prepared can be analyzed to determine the *A2M* genotype.

20 In a preferred embodiment of the invention, the sample(s) is genotyped for the *A2M-2* allele. In another preferred embodiment of the invention, the sample(s) is genotyped for the *A2M-G* allele.

In one embodiment of the invention, the genotype analysis is done utilizing RFLP analysis. RFLP analysis is a procedure well known in the art for identifying polymorphisms in DNA. After

25 restriction endonuclease digestion with one or more restriction enzymes, the DNA sample(s) is analyzed for a polymorphic variation. This step is usually accomplished by separation of the DNA fragments by size, a procedure which permits the determination of the presence of unique polymorphic fragments in one or more of the DNA samples. The fragments may be separated by any physical or biochemical means including gel electrophoresis, capillary electrophoresis,

30 chromatography (including sizing, affinity and immunochromatography), density gradient centrifugation and immunoabsorption. For carrying out the present invention, separation of DNA fragments by gel electrophoresis is particularly preferred, as it provides a rapid and highly reproducible means of sensitive separation of a multitude of DNA fragments, and permits direct,

5 simultaneous comparison of the fragments in several samples of DNA, or samples of DNA from a first and a second individual.

Gel electrophoresis is typically performed on agarose or polyacrylamide gels according to standard protocols. Samples are loaded onto the gels, usually with samples containing DNA fragments prepared from different sources of genomic DNA being loaded into adjacent lanes of the gel to facilitate subsequent comparison. Reference markers of known sizes may be used to facilitate the comparison of samples. Following electrophoretic separation, DNA fragments may be visualized and identified by a variety of techniques that are routine to those of ordinary skill in the art, such as ethidium bromide staining or autoradiography. One can then examine the samples for identifying polymorphic fragment patterns or for the presence of one or more unique bands.

A preferred embodiment of the invention is to utilize a restriction endonuclease specific for a restriction site created or deleted due to the mutation (substituting G for A) found in the *A2M-G* allele, and size fractionation by agarose or polyacrylamide gel electrophoresis to test for the occurrence of the *A2M-G* allele. A more preferred embodiment utilizes the restriction enzyme MboI for RFLP analysis to test for the occurrence of the *A2M-G* allele.

Another preferred embodiment of the invention is to analyze the sample for the occurrence of the *A2M-2* allele by digestion with a restriction endonuclease specific for a restriction site that has been created or deleted due to the pentanucleotide deletion found in the *A2M-2* allele, followed by size fractionation by agarose or polyacrylamide gel electrophoresis. One of ordinary skill in the art will be able to select an appropriate restriction endonuclease(s) for use in RFLP analysis by analyzing the site of the *A2M-2* pentanucleotide deletion. This can be done by determining whether the *A2M-2* allele contains a restriction site not found in the wild type *A2M* allele (*A2M-1*), or whether a restriction site found in the *A2M-1* allele in the region of the deletion mutation has been lost. Restriction sites for commercially available restriction endonucleases are described in Table 3.1.1 in Ausubel, F.M. *et al.*, eds., *Current Protocols in Molecular Biology*: John Wiley and Sons, Inc. (1997).

In a preferred embodiment of the invention, the DNA is amplified prior to the genotyping analysis. General methods for amplification and analysis of the DNA fragments are well-known to one of ordinary skill in the art (*see, e.g.*, U.S. Pat. Nos. 4,683,195; 4,683,202; and 4,800,159;

5 Innis, M.A., *et al.*, eds., *PCR Protocols: A Guide to Methods and Applications*, San Diego, California: Academic Press, Inc. (1990); Griffin, H.G., Griffin, A.M., eds., *PCR Technology: Current Innovations*, Boca Raton, Florida: CRC Press (1994) and Ausubel, F.M. *et al.*, eds., Chapter 15 in *Current Protocols in Molecular Biology*: John Wiley and Sons, Inc. (1997)).
10 Typically, these methods comprise contacting the DNA sample with a thermostable DNA polymerase in the presence of one or more primer sequences and amplifying the DNA sample to generate a collection of amplified DNA fragments, preferably by PCR or an equivalent automated amplification technique.

Based on the sequence of the alleles provided herein, PCR primers are constructed that are complementary to either (1) the region of *A2M* encompassing the site for the pentanucleotide
15 deletion found in the *A2M-2* allele, or (2) the region of *A2M* encompassing amino acid 1000 of the α_2 M protein. A primer consists of a consecutive sequence of polynucleotides complementary to any region in the allele encompassing the position of interest. The size of these PCR primers range anywhere from five bases to hundreds of bases. However, the preferred size of a primer is in the range from 10 to 40 bases, most preferably from 15 to 32 bases. As the size of the primer
20 decreases so does the specificity of the primer for the targeted region. Hence, even though a primer which is less than five bases long will bind to the targeted region, it also has an increased chance of binding to other regions of the template polynucleotide which are not in the targeted region. Conversely, a larger primer provides for greater specificity, however, it becomes quite cumbersome to make and manipulate a very large fragment. Nevertheless, when necessary, large
25 fragments are employed in the method of the present invention.

A more preferred method of the invention for amplifying the DNA is to utilize primers that flank the site of the pentanucleotide deletion mutation present in the *A2M-2* allele and provide a product that contains the sequence encoding exon 18 of α_2 M. More preferably, the invention utilizes primer 5' CTT TCC TTG ATG ACC CAA GCG CC 3' (SEQ ID NO: 1) and primer 5'
30 GTT GAA AAT AGT CAG CGA CCT C 3' (SEQ ID NO: 2) (Matthijs, G., and Marynen, P., *Nucl. Acids Res.* 19(18):5102 (1991)) during the amplification step.

Another preferred method of the invention for amplifying the DNA is to utilize primers that flank the sequence encoding amino acid 1000 of the α_2 M protein and provide a product that contains the sequence encoding said amino acid. More preferably, the invention utilizes the sense

5 primer C23: 5' ATC CCT GAA ACT GCC ACC AA 3' (SEQ ID NO: 3) and antisense primer AS24 : 5' GTA ACT GAA ACC TAC TGG AA 3' (SEQ ID NO: 4) during the amplification step.

Following amplification by the methods of the present invention, the amplified DNA fragments may be analyzed to identify a specific polymorphism. The analysis can be done by
10 determining the polynucleotide sequence of the amplified fragments using procedures well known in the art. Several methods of sequencing DNA are described by Ausubel, F.M. *et al.*, eds., in Chapter 7 in *Current Protocols in Molecular Biology*: John Wiley and Sons, Inc. (1997) (*see also*, U.S. Patent Nos. 4,962,022 and 5,498,523, which are directed to methods of DNA sequencing). This method of analysis is preferred when the amplified fragment contains the site
15 of the deletion mutation found in the *A2M-2* allele.

In another embodiment, the analysis can be done by RFLP analysis, as was previously described herein. This method of analysis is preferred when the fragment amplified encodes amino acid 1000 of the α_2 M protein.

In yet another embodiment, analysis of the amplified sample can be done by Single-Strand
20 conformation polymorphism (SSCP) analysis. SSCP analysis is a procedure well known in the art for identifying polymorphisms in DNA (*see, e.g.*, Orita, M., *et al.*, *Proc. Natl. Acad. Sci. USA* 86:2766-2770 (1989); Teschauer, W., *et al.*, *Eur. J. Clin. Chem. Clin. Biochem.* 34:125-131 (1996); and Paccoud, B., *et al.*, *Nucl. Acids Res.* 26(9):2245-2246 (1998)). After denaturation of a double stranded DNA sample(s), the single stranded DNA is analyzed for a polymorphic
25 variation. This step is usually accomplished by separation of the single stranded DNA using electrophoresis, a procedure which permits the determination of the presence of unique polymorphic conformations in the DNA sample(s). The strands may be separated on a non-denaturing polyacrylamide gel. Gel electrophoresis and visualization of the DNA may be performed as described immediately herein.

30 In an additional embodiment, the amplified sample can be analyzed directly by size fractionation if the sequence of interest contains an insertion or deletion relative to the control DNA.

An alternative embodiment of the invention utilizes protein isotyping to determine the *A2M* genotype. The α_2 M variant found in the *A2M-2* genotype is predicted to lack exon 18 due

5 to the loss of nucleotides in the splice acceptor site of exon 18. These nucleotides are believed to be necessary for effective splicing of α_2 M. In the *A2M-G* genotype, Ile¹⁰⁰⁰ (ATC) is replaced by Val¹⁰⁰⁰ (GTC). Antibodies specific to each of these variant forms of α_2 M can be utilized to screen individuals for the occurrence of these variants and thus for the occurrence of the *A2M-2* or *A2M-G* alleles.

10 The term "antibody" refers both to monoclonal antibodies which have a substantially homogeneous population and to polyclonal antibodies which have heterogeneous populations. Polyclonal antibodies are derived from the antisera of immunized animals, and monoclonal antibodies to specific α_2 M variants may be obtained by methods known in the art (Kohler and Milstein, *Nature* 256:495-497, 1975). Such antibodies may be of any immunoglobulin class
15 including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The term "antibody" is meant to include both intact molecules as well as fragments thereof, such as Fab and F(ab')₂, which are capable of binding antigen. The antibodies are tested for their ability to distinguish the α_2 M-2 from the α_2 M-1 proteins, or the Val¹⁰⁰⁰ from the Ile¹⁰⁰⁰ proteins by a standard immunoassay method such as ELISA (Chapter 11 in *Current Protocols In Molecular Biology*: Green Publishing
20 Associates and Wiley-Interscience (1991)), using recombinantly expressed α_2 M.

Once tested, the antibodies may be used to detect the presence of the α_2 M-2 variant or the α_2 M Val¹⁰⁰⁰ variant. These antibodies may be used in immunoassay techniques well known in the relevant arts, such as Western Blot or ELISA. Thus, in one embodiment, an antibody specific for the α_2 M-2 variant is used to assess *A2M* genotype through protein isotyping. In a preferred
25 method of protein isotyping, a second antibody, which is specific for α_2 M-1 is used as a control. Yet another embodiment of the invention relates to the use of an antibody specific for the Val¹⁰⁰⁰ variant to assess the *A2M-G* genotype through protein isotyping. In a preferred method of protein isotyping, a second antibody, which is specific for the Ile¹⁰⁰⁰ protein, is used as a control. For the protein isotyping methods of the invention, preferably, protein is isolated from an individual and
30 screened with the specific antibodies.

An alternative method of protein isotyping the *A2M-2* genotype is to assay for the α_2 M-2 variant using an α_2 M electrophoretic-mobility assay, which consists of analyzing the electrophoretic mobility of α_2 M under non-denaturing conditions after incubation with a protease, or other reagent capable of converting α_2 M to the fast form (Barret, A. J., *et al.*, *Biochem. J.* 181:

5 401-418 (1979); Bowen, M. E., and Gettins, P. W., *J. Biol. Chem.* 273:1825-1831 (1998)). α_2 M can exist in two forms easily distinguishable by mobility in gel electrophoresis (Barret, A. J., *et al.*, *Biochem. J.* 181: 401-418 (1979)). The difference in mobility is due to a conformational change. α_2 M has a tetrameric structure that undergoes a conformational change upon proteolytic cleavage near the center of the bait domain by a protease (Borth, W. *FASEB* 6: 3345-3353 (1992)). This conformational change traps the protease ((Borth, W. *FASEB* 6: 3345-3353 (1992))), and results in an increase in electrophoretic mobility on poly-acrylamide gels run under non-denaturing conditions (this form is referred to as the "fast form" of α_2 M) (Barret, A. J., *et al.*, *Biochem. J.* 181: 401-418 (1979); Bowen, M. E., and Gettins, P. W., *J. Biol. Chem.* 273:1825-1831 (1998)). This "slow to fast" conversion is used as the basis for an assay for this conformational change, and the two different α_2 M conformations are referred to as the slow and fast forms (Bowen, M. E., and Gettins, P. W., *J. Biol. Chem.* 273:1825-1831 (1998)). This assay is referred to herein as the α_2 M electrophoretic mobility assay.

Because the α_2 M-2 variant is expected to lack exon 18, the second of the two exons of the bait domain, it is also expected that this variant would be unable to undergo the conformational change that increases electrophoretic mobility converting it from the slow form to the fast form of α_2 M. Thus, the α_2 M-2 variant can be identified using the α_2 M electrophoretic mobility assay.

The α_2 M electrophoretic mobility assay and methods of purifying α_2 M from serum are described by Barret *et al.* in Barret, A. J., *et al.*, *Biochem. J.* 181: 401-418 (1979), and by Bowen *et al.* in Bowen, M. E., *et al.*, *Arch. Biochem. Biophys.* 337:191-201 (1997), and in Bowen, M. E., and Gettins, P. W., *J. Biol. Chem.* 273:1825-1831 (1998). Reagents for use in the conversion of α_2 M to the fast form include proteases, such as trypsin, papain, thermolysin or chymotrypsin; ammonia or low-molecular-weight ammonium analogues, such as $(\text{NH}_4)_2\text{SO}_4$, methylammonium chloride and ethylammonium chloride; and salts of certain amines such as 1,3-diaminopropane, ethanolamine and methylamine (Bowen, M. E., and Gettins, P. W., *J. Biol. Chem.* 273:1825-1831 (1998)). After incubation with one of these reagents, preferably a protease, the α_2 M sample may be run on polyacrylamide gel under nondenaturing conditions, such as those described in Bowen, M. E., *et al.*, *Arch. Biochem. Biophys.* 337:191-201 (1997). The α_2 M sample may be detected by methods well known in the art such as by radiolabelling the protease used, or by Western Blot

5 using anti- α_2 M antibodies. The α_2 M-2 variant, and/or the fast form of α_2 M-1 may be used as controls for comparison of electrophoretic mobility with the sample being analyzed.

The invention also provides kits for use in the identification, analysis and typing of polymorphic DNA according to the present methods. Kits according to the present invention may comprise a carrying means being compartmentalized to receive in close confinement therein one
10 or more containers such as vials, tubes, bottles and the like. Each of such containers may comprise components or a mixture of components needed to perform genotyping analysis.

A kit for genotyping DNA can comprise a number of containers. In one example of such a kit, a first such container can contain a substantially purified sample of the primers with the sequences of SEQ ID NOs 1 and 2 described above. A second such container can contain α_2 M-2
15 DNA. A third such container may contain α_2 M-1 DNA. In another example of a kit for genotyping DNA, a first such container can contain a substantially purified sample of the primers with the sequences of SEQ ID NOs 3 and 4 described above. A second such container can contain α_2 M Val¹⁰⁰⁰ DNA. A third such container may contain α_2 M Ile¹⁰⁰⁰ DNA.

In another embodiment, the invention provides a kit for *A2M* genotyping by means of
20 protein isotyping. Such a kit can include one or more containers. In an example of such a kit, a first such container may contain an antibody specific to the α_2 M-2 protein. A second such container may contain an antibody specific to the α_2 M-1 protein. A third such container may contain a substantially purified sample of the α_2 M-2 variant, or fragment thereof. A fourth such container may contain a substantially purified sample of α_2 M-1 protein, or fragment thereof. In
25 another example of a kit for *A2M* genotyping by means of protein isotyping, a first such container may contain an antibody specific to the α_2 M Val¹⁰⁰⁰ variant. A second such container may contain an antibody specific to α_2 M Ile¹⁰⁰⁰ protein. A third such container may contain a substantially purified sample of the α_2 M Val¹⁰⁰⁰ variant, or fragment thereof. A fourth such container may contain a substantially purified sample of α_2 M Ile¹⁰⁰⁰ protein, or fragment thereof. In yet another
30 example of a kit for *A2M* genotyping by means of protein isotyping, a first container may contain a protease or other reagent for use in conversion of an α_2 M to the fast form. A second such container may contain a substantially purified sample of the α_2 M-2 variant. A third such container may contain a substantially purified sample of the fast form of α_2 M-1. Other containers may contain reagents for use in the purification of α_2 M or for use in electrophoresis.

5 It will be readily apparent to those skilled in the relevant arts that other suitable modifications and adaptations to the methods and applications described herein are obvious and may be made without departing from the scope of the invention or any embodiment thereof. Having now described the present invention in detail, the same will be more clearly understood
10 by reference to the following examples, which are included herewith for purposes of illustration only and are not intended to be limiting of the invention.

EXAMPLES

15 To identify novel AD genes, the inventors employed a candidate gene strategy focusing on genes for known LRP ligands other than apoE and APP (Kounnas, M., *et al.*, *Cell* 82:331-340 (1995)). α_2 M is a major LRP ligand and major serum pan-protease inhibitor which inhibits all four classes of proteases (Borth, W., *FASEB* 6:3345-3353 (1992)). In brain, *A2M* is upregulated during injury along with its major receptor, the low density lipoprotein receptor-related protein (LRP) (Strickland, D. K., *et al.*, *J. Biol. Chem.* 265:17401-17404 (1990); Businaro, R., *et al.*,
20 *Exp. Brain Res.* 88:213-218 (1992); Lopes, M. B., *et al.*, *FEBS Lett.* 338: 301-305 (1993)). α_2 M has been localized to senile plaques (SP) in AD (Bauer, J., *et al.*, *FEBS* 285:111-114 (1991)) and tightly binds the β AP peptide, the major component of β -amyloid in SP and cerebral blood vessel deposits (Du, Y., *et al.*, *J. Neurochem.* 69:299-305 (1997) and Hughes, S. R., *et al.*, *Proc. Natl. Acad. Sci. (USA)* 95:3275-3280 (1998)). α_2 M has also been shown to attenuate the fibrillogenesis
25 and neurotoxicity of β AP (Hughes, S. R., *et al.*, *Proc. Natl. Acad. Sci. (USA)* 95:3275-3280 (1998) and Du, Y., *et al.*, *J. Neurochem.* 70:1182-1188 (1998)) and to mediate both its degradation (Qiu *et al.*, *J. Biol. Chem.* 271:8443-8451 (1996)) and clearance via endocytosis through LRP (Narita, M., *et al.*, *J. Neurochem.* 69:1904-1911 (1997)). In view of these findings
30 implicating α_2 M in AD, the inventors tested for a genetic association between *A2M* and AD.

During the course of this study, chromosome 12 linkage to AD was reported as part of
a genome scan (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1244 (1997)). Since *A2M* maps within 30 cM of the implicated chromosome 12 markers (Krauter, K., *et al.*, *Nature* 377:321-333 (1995)), the inventors attempted to confirm this result using the National Institute of Mental Health (NIMH) Genetics Initiative AD sample (Blackner, D., *et al.*, *Neurology* 48:139-

147 (1997)) and to determine whether a genetic association of *A2M* with AD may account for the previously published chromosome 12 linkage.

Example 1

Pentanucleotide Deletion Allele

A. Methods

1. Sample

To test for a link between *A2M* and AD, the inventors used the National Institute of Mental Health (NIMH) Genetics Initiative AD sample, a large sample of affected sibling pairs and other small families with AD (Blacker, D., *et al.*, in *Neurology* 48:139-147 (1997)). Participants were recruited from local memory disorder clinics, nursing homes, and the surrounding communities with the only requirement for inclusion in the sample being that each family include at least two living blood relatives with memory problems. They were evaluated following a standardized protocol (Blacker, D., *et al.*, *Arch. Neurol.* 51:1198-1204 (1994)) to assure that they met NINCDS/ADRDA criteria for Probable AD (or in the case of secondary probands, Possible AD) (McKhann, G., *et al.*, *Neurology* 34:939-944 (1984)), or research pathological criteria for Definite AD (Khachaturian, Z., *Arch. Neurol.* 42:1105 (1985)). Among the affected individuals, 142 (22.2%) had autopsy confirmation of the diagnosis of AD. Unaffected relatives, generally siblings, were included when they were available and willing to participate.

There were a total of 239 unaffected subjects from 131 families (45.6%). An additional 22 study subjects with blood available who had unclear phenotypes were considered phenotype unknown, as were 5 unaffected subjects with unknown ages, and 19 unaffected subjects below 50 years of age (primarily children of affected participants). There were a total of 639 individuals affected with AD, from 286 families. The majority of the affected individuals were sibling pairs (202 families, 71%), but there were 46 larger sibships (16%), and 38 families with other structures (13%; e.g., parent-child, first cousin, avuncular, extended). All subjects (or, for significantly cognitively impaired individuals, their legal guardian or care-giver with power of attorney) gave informed consent.

5 The full sample was used in the descriptive statistics for genotype counts and allele frequencies, for the analyses of age of onset in affected individuals, and for all of the genetic linkage analyses (except ASPEX, which uses sibships only). However, because the Mantel-Haenzel test, conditional logistic regression, and Sibship Disequilibrium Test depend on comparisons of closely related affected and unaffected individuals, these were performed on a
10 subsample including all families in which there was at least one affected and at least one unaffected sibling with *A2M* data available: 104 families with 217 affected and 181 unaffected siblings.

 In order to avoid examining very early onset AD, which appears to have a distinct genetic etiology (Blacker, D., & Tanzi, R. E., *Arch Neurol* 55:294-296 (1998)), only those families in
15 which all examined affected individuals experienced the onset of AD at age 50 or later were included. Although LOAD is conventionally identified based on onset after age 60, families with onsets between 50 and 60 were included because onset in this decade is only partly explained by the known AD genes. Age of onset was determined based on an interview with a knowledgeable informant and review of medical records.

20 2. *Amplification and Genotyping of A2M, APOE, and Chromosome 12 Markers*

 The *A2M* exon 18 splice acceptor pentanucleotide deletion (the *A2M-2* allele) was manually genotyped according to the protocol described in Matthijs *et al.* (Matthijs, G., &
25 Marynen, P., *Nuc. Acid. Res.* 19:5102 (1991)). A 326 bp fragment was amplified by PCR using primers having the sequences described in SEQ ID NOs 1 and 2. The amplified fragment was then sequenced using 33P in a 96-well format on 6% denaturing polyacrylamide gels. The presence of the deletion resulted in a 5 bp smaller amplicon. *APOE* was genotyped as described
in Blacker *et al.* (Blacker, D., *et al.*, *Neurology* 48:139-147 (1997)). Chromosome 12 markers
30 were genotyped using primers that were either provided from the Weber 8 set (Research Genetics) or were custom synthesized (BRL).

 Manual genotyping was carried out using a 96-well microtiter dish format. Three to 10 nanograms, of human DNA was mixed with a reaction buffer, deoxynucleotide mix (e.g. for a poly-[dGdT] STR, the final concentration is 200 mM each of dATP, dCTP and dTTP; and 2 mM

5 dGTP), 1 mCi alpha-32PdGTP or 33P-dGTP, 15 pM of each flanking primer and .25 units of Taq polymerase in a total volume of 10 ml. The reactions were denatured at 94°C for 4 min, followed by 25-30 cycles of 1 min denaturing at 94°C, 0.5-1 min annealing (variable temperature, usually 55-65°C) and extension for 1 min at 72°C. Forty-eight (48) experimental and two control (for standardization of size) samples were loaded on a gel at one time, thereby increasing the amount of information per gel. Whenever possible (e.g., if marker background was sufficiently low) multiple markers (two to four markers) were multiplexed, or were temporally staggered (30-45 minutes) two to three mm on a single gel. Allele sizes for CEPH individuals 1331-01 and 1331-02 were used as standards. In the rare event that no standards were available for a marker, an initial gel was run which included a sequencing ladder to determine allele sizes in these individuals. Two
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075
10080
10085
10090
10095
10100
10105
10110
10115
10120
10125
10130
10135
10

5 were performed in the SAS statistical analysis package (SAS Institute, SAS Program Guide, Version 6, Cary, NC (1989)).

For all types of analysis, allele frequencies were computed from the data, but rare alleles were adjusted up to a frequency of 0.01 (with a compensatory small decrease in the frequency of the most common alleles) in order to minimize the possibility of a false positive result. All analyses were repeated using the uncorrected frequencies.

B. Results

1. Genotype and Allele Frequencies

15 For descriptive purposes, *A2M* genotype counts and allele frequencies were examined in affected and unaffected subjects in study families. Unaffected individuals in AD families are not genetically independent of their affected relatives, of course, and thus would be expected to show higher frequencies of AD-associated alleles compared to the general population. However, given an increased risk of AD with a given allele, its frequency would be expected to be higher among affected individuals than among their unaffected relatives. However, since these frequencies are pooled across families, they are neither as accurate nor as powerful an indicator of genetic association as the SDT.

25 *A2M* genotype counts and allele frequencies for *A2M-2* are reported separately for primary and secondary probands, with primary probands serving as the primary subject population, and secondary probands as a confirmation sample. Allele frequencies in the probands were compared to those for unaffected individuals based on the oldest unaffected individual from each of the 105 families in which one or more unaffected subjects with *A2M* data was available. In addition, the analyses were repeated using an unaffected sample that had passed through a majority of the age of risk, the "stringent" unaffecteds, those who were at least as old as the age of onset of the latest-onsetting affected family member, again selecting the oldest such individual in each family. Because age of onset is correlated in families (Farrer, L. A., *et al.*, *Neurology* 40:395-403 (1990)), using onset ages in the subjects' own families is preferable to setting an arbitrary cutoff.

35 The applicants initially determined genotype counts and allele frequencies for the bi-allelic 5' exon 18 splice acceptor deletion polymorphism in *A2M* (Matthijs, G., Marynen, P., *Nuc. Acid.*

5 *Res. 19:5102 (1991))* in primary probands, secondary probands, unaffected individuals (oldest in family), and "stringent" unaffecteds, (those who have reached the onset age of the latest-onsetting affected, again using the oldest such individual), stratified on individual *APOE* dose (see Tables 1 and 2). The combined genotype frequencies for the possession of 1 or 2 *A2M-2* alleles was higher in the primary and secondary probands (29.5% and 31.9%) than in the oldest unaffecteds (22.9%) and oldest stringent unaffected (19.3 %; Table 1). The *A2M-2* allele frequencies for primary and secondary probands; were similar (16.4% and 17.7%) and both were lower than those observed in unaffecteds individuals (12.9% for oldest unaffecteds and 10.5% for oldest stringent unaffecteds; Table 2). The effect was particularly marked among the *APOE-ε4* zero-dose individuals (15.0% for primary probands and 19.4% for secondary probands vs. 4.3% for oldest unaffecteds and 3.7% for oldest stringent unaffected).

20

25

Table 1: <i>A2M</i> Genotypes Stratified on Individual <i>APOE-ε4</i> Dose												
	Primary Probands			Secondary Probands			Oldest Unaffecteds			Oldest Stringent Unaffecteds*		
<i>A2M</i> genotype	1/1	1/2	2/2	1/1	1/2	2/2	1/1	1/2	2/2	1/1	1/2	2/2
<i>APOE-ε4</i> dose												
0	59	18	3	62	26	5	42	4	0	25	2	0
1	100	39	3	89	37	2	33	12	2	22	7	1
2	42	19	3	43	18	3	6	5	1	3	2	0
Total sample	201	76	9	194	81	10	81	21	3	50	11	1
Percent	70.3	26.4	3.1	68.1	28.4	3.5	77.1	20.0	2.9	80.6	17.7	1.6
*Stringent unaffecteds are those at least as old as the latest onsetting affected subject in the family.												

Table 2: *A2M-2* Allele Frequencies Stratified on Individual *APOE-ε4* Dose

	Primary Probands		Secondary Probands		Oldest Unaffecteds		Oldest Stringent Unaffecteds*	
	freq	n ^s	freq	n ^s	freq	n ^s	freq	n ^s
<i>APOE-ε4</i> dose								
0	.150	80	.194	93	.043	46	.037	27
1	.158	142	.160	128	.170	47	.150	30
2	.195	64	.188	64	.292	12	.200	5
Total	.164	286	.177	285	.129	105	.105	62

*Stringent unaffected are those at least as old as the latest onsetting affected subject in the family.

^sn refers to the number of subjects on which the frequency is based, and is thus half the number of alleles.

2. Mantel-Haenzel Odds Ratios and Conditional Logistic Regression

Mantel-Haenzel odds ratios were calculated for the odds of being affected given the possession of at least one *A2M-2* allele. These analyses were performed stratified on family using n-to-m matching, so all members of a sibship could be used and intercorrelations among siblings could be taken into account. Spielman and Ewens (Spielman, R. S., and Ewens, W. J., *Am. J. Hum. Genet.* 62:450-458 (1998)) have recently suggested the use of a similar analysis to test for linkage. The analyses were performed first using all unaffected siblings, and then only the stringent unaffected siblings.

Conditional logistic regression was used to control the Mantel-Haenzel odds ratio for the effect of *APOE-ε4* on AD risk. Here, the outcome is disease status of each sibling, conditioning on family using an n-to-m matching paradigm, and including *APOE-ε4/ε4* homozygosity as a covariate, along with a term for the interaction between *APOE-ε4* and *A2M-2*. Like the Mantel-Haenzel odds ratio, conditional logistic regression is a standard method for analysis of data from matched sets, and can control for clustering of genotypes within families of arbitrary size. These analyses were performed using the PHREG procedure in SAS (SAS Institute, SAS Program Guide, Version 6, Cary, NC (1989)). These analyses were repeated using only the "stringent"

5 unaffected siblings (those who were as least as old as the onset age of the oldest-onsetting affected sibling) in order to minimize the effect of misclassification of unaffected siblings. These analyses were also performed coding *APOE-ε4* as gene dosage, and including a term for the possession of an *APOE-2* allele, previously shown to decrease disease risk (Corder, E. H., *et al.*, *Nat Genet.* 7:180-184 (1994); Farrer, L. A., *et al.*, *JAMA* 278:1349-1356 (1997)).

10 The Mantel-Haenzel odds ratio (Table 3) for the risk of being affected as a function of carrying at least one *A2M-2* allele was 3.56 (95% CI = {1.80, 7.03}; $p = .0003$). In comparison, this same sample set yielded a similar odds ratio for the risk of being affected as a function of carrying two copies of *APOE-ε4* (3.54 [95% CI = {1.76, 7.12}]; $p = .0004$). Thus, risk due to inheriting at least one *A2M-2* allele was comparable to that conferred by *APOE-ε4* homozygote status. When this analysis was restricted to stringent unaffecteds, a much smaller sample, the
15 Mantel-Haenzel odds ratio for *A2M-2* remained significant (2.27 [95% CI = {1.19, 6.23}]; $p = .015$). Conditional logistic regression analyses, which allow for the calculation of Mantel-Haenzel odds ratios adjusted for the effect of *APOE-ε4* on AD risk are also shown in Table 3, first for all affected and unaffected siblings, and then for all affected and stringent unaffected siblings. Of
20 note, comparable and significant magnitudes of risk were conferred by the possession of one or more *A2M-2* alleles (Model 1) or by *APOE-ε4* homozygosity (Model 2) and the magnitude of that risk changed little when both genes were included in the model (Model 3). There was also no evidence of an interaction between the two genes on disease risk (Model 4). The results for analyses using only the affecteds and stringent unaffecteds were similar, but less significant, as
25 would be expected given the smaller sample size. Additional analyses were performed (not shown) coding *APOE-ε4* "dose" as an ordinal variable and including *A-POE-2* in the model, with no appreciable difference in the results.

Table 3: Conditional logistic regression for the effect of *A2M* and *APOE-ε4* on risk for AD

All Affecteds and Unaffecteds				
Model	Variables	est. OR	95% CI	p-value
1	<i>A2M</i> any 2	3.56	(1.80, 7.03)	.0003
2	<i>APOE-ε4/ε4</i>	3.54	(1.76, 7.12)	.0004
3	<i>A2M</i> any 2 <i>APOE-ε4/ε4</i>	3.45 3.45	(1.71, 6.94) (1.67, 7.10)	.0005 .0008
4	<i>A2M</i> any 2 <i>APOE-ε4/ε4</i> interaction	3.40 3.39 1.07	(1.57, 7.35) (1.51, 7.64) (0.25, 4.46)	.0018 .0032 .932
All Affecteds and Stringent* Unaffecteds				
1	<i>A2M</i> any 2	2.72	(1.19, 6.23)	.018
2	<i>APOE-ε4/ε4</i>	3.90	(1.62, 9.40)	.0025
3	<i>A2M</i> any 2 <i>APOE-ε4/ε4</i>	2.85 4.10	(1.21, 6.70) (1.66, 10.12)	.017 .0023
4	<i>A2M</i> any 2 <i>APOE-ε4/ε4</i> interaction	2.48 3.61 1.78	(0.96, 6.39) (1.36, 9.58) (0.29, 10.9)	.061 .0099 .533

* Stringent unaffecteds are those at least as old as the latest onsetting affected subject in the family.

3. Sibship Disequilibrium Test

The Sibship Disequilibrium Test (SDI) (Horvath and Laird, under revision, *Am. J. Hum. Genet.*) is a non-parametric sign test developed for use with sibling pedigree data that compares the average number of candidate alleles between affected and unaffected siblings. The SDT is similar to the Sib-TDT, a recently developed test that also does not require parental data (Spielman, R. S., and Ewens, W. J., *Am. J. Hum. Genet. (Suppl.)* 53:363 (1993)), but has the advantage of being able to detect association in sibships of an arbitrary size. Like the TDT, Sib-TDT, and other family-based association tests, the SDT offers the advantage of not being susceptible to errors due to admixture. Another advantage of these methods is that misclassification of affection status (*e.g.*, due to the unaffected siblings not having passed through the age of risk) decreases the power of the test, but does not lead to invalid results. The SDT can

5 test both linkage and linkage disequilibrium; it can only detect linkage disequilibrium in the presence of linkage, hence there is no confounding due to admixture. The null hypothesis of the SDT is that $\Theta = 1/2$ (no linkage) or $\delta = 0$ (no disequilibrium), i.e., $H_0: \delta(\Theta - 1/2) = 0$. The STD program (for several platforms) and documentation may be found at <ftp://sph70-57.harvard.edu/XDT/>.

10 Because the SDT does not require parental data, and can use all information from sibships of arbitrary size, it is well-suited to the analysis of the NIMH AD data. Before using it to detect novel AD genes, the SDT was validated with the known AD gene *APOE-ε4* in the sample. In an examination of 150 sibships with 286 affected and 242 unaffected individuals from the sample (n.b., the number of sibships is higher than that for the *A2M* analyses reported here because a
15 greater number of families have been typed for *APOE*), the SDT was able to detect not only the deleterious *APOE-ε4* effect but also the more difficult to detect *APOE-2* protective effect (Farrer, L. A., et al., *JAMA* 278:1349-1356 (1997); Corder, E. H., et al., *Nature Genet.* 7:180-184 (1994)) not previously detected in these data (Blacker, D., et al., *Neurology* 48:139-147 (1997), and see Wilcox et al., in preparation).

20 The primary analysis of the association of *A2M-2* with AD examined the probability of passing along an *A2M-2* allele as a function of affection status. In order to increase the likelihood of correct classification of unaffected status, the analyses were repeated including only "stringent" unaffected siblings, those who were at least as old as the latest onset affected siblings, a sample of 60 families. In addition, in order to assess whether the effect differed in different *APOE*
25 genotypes persisted in individuals with similar *APOE* genotypes, the analyses were repeated within strata defined by matching affected and unaffected siblings for *APOE-ε4* gene dose. There were 18 *APOE-ε4* zero-dose sibships, 21 *APOE-ε4* one-dose sibships, and 11 *APOE-ε4* two-dose sibships.

30 Analysis by the SDT revealed that *A2M-2* confers significant risk for AD in the total sample ($Z = 4.74$; $p = 0.00009$). Once again the magnitude of the association was comparable to that observed in individuals possessing two copies of *APOE-ε4* ($Z = 4.49$; $p = 0.00006$). The results of the SDT are shown in Table 4, including Z values (indicating the magnitude and direction of the effect), the exact p -values, the total number of sibships involved, and the number of sibships with a difference in the mean number of candidate alleles (*A2M-2* or *APOE-ε4*)

5 between affected and unaffected siblings (*i.e.*, the effective sample size). This effect persisted when only the stringent unaffected siblings were used, and within the subset of sibships concordant for zero or one "dose" of *APOE-ε4* (data not shown). To provide further validation of the SDT, the Sibling TDT (Spielman, R. S. and Ewens, W. J., *Am. J. Hum. Genet.* 62:450-458 (1998)) (Sib-TDT) was applied. The Sib-TDT also revealed significant association between *A2M* and AD ($Z = 3.61$, $p = 0.0002$).

Table 4: Sibling disequilibrium test* results for association between the *A2M-2* allele and AD

Sample	Number of sibships	Effective sample size [#]	Z ^s	p
<i>A2M-2</i> : Total sample	104	40	4.74	0.00009
<i>A2M-2</i> : Sibships including a stringent unaffected [@]	60	27	3.47	0.0059
<i>APOE-ε4</i> : Total sample	104	47	4.49	0.00006

* Horvath and Laird, under revision, AJHG.

[#] The effective sample size is the number of sibships with a difference in the mean number of candidate alleles (*A2M-2*, *APOE-ε4*) between affected and unaffected siblings.

^s Z values indicate the magnitude and direction of the effect.

[@] Stringent unaffecteds are those at least as old as their oldest onsetting affected sibling.

4. Age of Onset Based Techniques

30 In order to see if *A2M* effects appeared to operate via changes in age of onset, affected individuals were examined according to *A2M* genotype, stratifying on or controlling for the powerful effect of *APOE-ε4*. First, this was examined graphically using Kaplan Meier curves (data not shown) including all affected and unaffected individuals, first stratifying on *A2M* genotype alone, and then on *A2M-2* carrier status and *APOE-ε4* dose. Second, the mean ages of onset of primary and secondary probands were compared by *A2M* genotype overall, and stratified on *APOE-ε4* gene dose. Third, analysis of variance (performed separately for primary and secondary probands) was used, including first only *A2M* genotype (defined as any 2 vs. none),

then only APOE genotype (defined as *APOE-ε4* gene dose or *APOE-ε4/ε4* vs. not), then both, and then both plus an interaction term.

Although *A2M-2* appeared to increase risk for AD, unlike *APOE-ε4*, it did not appear to decrease age of onset of AD. This is shown graphically in Kaplan Meier curves of the sample as a whole, stratified first on *A2M* genotype alone and then on both the presence of an *A2M-2* allele and *APOE-ε4* dose. These graphs clearly show that inheritance of the *APOE-ε4/ε4* genotype lowers age of onset, irrespective of *A2M* genotype. However, no difference in age of onset was conferred by *A2M* genotype for the overall sample or within any *APOE-ε4* dose stratum.

The mean ages of onset for primary and secondary probands for each *A2M* genotype overall, and stratified according to individual *APOE-ε4* dose showed no effect of *A2M* genotype on age of onset of AD (Table 5), but did show the expected decline in age of onset homozygous for *APOE-ε4*. Analysis of variance of primary (and secondary, performed separately) probands showed no effect of *A2M* genotype on age of onset, but did show a significant effect of the *APOE-ε4/ε4* genotype on age of onset.

Table 5: Age of onset of AD by *A2M* genotype, stratified on individual *APOE-ε4* dose

		Primary Probands			Secondary Probands		
	<i>A2M</i> Genotype	1/1	1/2	2/2	1/1	1/2	2/2
<i>APOE</i> dose							
0	n m (sd)	59 73.0(8.7)	18 76.2(10.0)	3 76.3(3.5)	62 74.7(8.2)	26 75.7(8.2)	5 76.4(7.2)
1	n m (sd)	100 71.7(7.4)	39 73.0(7.9)	3 73.0(7.9)	89 73.8(5.9)	37 73.0(6.0)	2 70.0(4.2)
2	n m (sd)	42 68.4(6.9)	19 62.8(6.0)	3 67.0(4.4)	43 65.6(6.7)	18 67.9(8.8)	3 65.0(7.9)
Total	n m (sd)	201 71.4(7.8)	76 71.7(9.5)	9 72.1(6.4)	194 72.3(7.3)	81 72.7(7.8)	10 71.7(8.2)

5. *Traditional Linkage Analysis and Related Techniques*

In order to determine if there was evidence for linkage in the *A2M* region, and in particular whether *A2M* might be related to the recent reports of linkage to the centromeric region of chromosome 12 (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)), a variety of genetic linkage analytic techniques were performed. For these analyses, all families were divided into "tiers" according to the criteria of Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)): Tier 1 (30 families) is comprised of families in which all affecteds were *APOE-ε4/ε4*; Tier 2 (131 families) is comprised of families not in Tier 1 in which all affecteds were *APOE-ε4* carriers; and Tier 3 (126 families) is comprised of families in which at least one affected did not carry an *APOE-ε4* allele. It should be noted that Pericak-Vance *et al.* found evidence for linkage to chromosome 12 markers only for Tier 3. All analyses were run on the entire sample and on each of these three tiers. In addition, because the analyses in Pericak-Vance *et al.* were confined to families with all onsets at 60 or above, and in which AD was evident in at least two generations, the analyses were repeated using the 259 families meeting these criteria (24 in Tier 1, 118 in Tier 2, and 117 in Tier 3).

The first technique used was conventional linkage analysis using two autosomal dominant disease models. The first model was an affecteds only analysis based on the model used by Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)): a gene frequency of .001 and a phenocopy rate of .05. The other main model was an age-curve model assuming a normal distribution of disease onset with a mean of 71.4 and sd of 8.7 (*n.b.*, these are the observed values in the Genetics Initiative sample, and are very similar to those used in Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997))), a disease gene frequency of .01, and a fixed phenocopy rate of .10 (*n.b.*, this value was used for the Probable AD cases [70.3 % of the subjects] and for the computation of partial penetrances for unaffected subjects; a penetrance of .05 was used for the Definite AD cases [22.2% of the subjects], and .14 for the Possible AD cases [7.5% of the subjects])). However, because these analyses were done in part in an attempt to replicate the findings of Pericak-Vance *et al.*, additional models (all using the normal onset distribution described above) considered consistent with AD genetics and prevalence were also tested, including the inventors' best approximation of the age curve model

5 used by Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)),
with a disease gene frequency (q) of .001 and a phenocopy rate (ϕ) of .05; $q = .01$ and $\phi = .05$; q
= .02 and $\phi = .05$; $q = .02$ and $\phi = .10$. All lod score analyses were performed in Fastlink
(Terwilliger, J. D., Ott, J., *Handbook of Human Genetic Linkage*. Baltimore: Johns Hopkins UP,
(1994)).

10 The second technique was multipoint non-parametric linkage analysis (Figures 2A-2D)
using the program GENEHUNTER (Kruglyak, L., *et al.*, *Am J Hum Genet* 58:1347-1363
(1996))(software available from Kruglyak *et al.* by anonymous ftp at ftp-genome.wi.mit.edu or
from their World Wide Web site <http://www-genome.wi.mit.edu/ftp/distribution/software/genehunter>)). a multipoint non-parametric linkage program accommodating full pedigrees, and
15 reporting non-parametric lod (NPL) scores.

The last technique was the multivariate sibship analysis package ASPEX (Affected Sibpair
Exclusion Mapping; <ftp://lahmed.stanford.edu/pub/aspe/>), which is based on allele sharing within
sibships. The analyses were performed using the Sib-Phase program, using fixed, allele
frequencies based on those observed in the data (see below) to estimate IBD probabilities when
20 parental information is missing, and using no dominance variance. In addition, exclusion analysis
was performed setting the ASPEX parameter "risk" (which is roughly equivalent to X , the
recurrence risk in relatives) at 2, a reasonable value for AD based on family data (Farrer, L. A.,
et al., *Neurology*, 40(3 Pt 1):395-403 (1990)).

For all types of analysis, allele frequencies were computed from the data, but rare alleles
25 were adjusted up to a frequency of .01 (with a compensatory small decrease in the frequency of
the most common alleles) in order to minimize the possibility of a false positive result. All
analyses were repeated using the uncorrected frequencies.

For the multipoint techniques (ASPEX and GENEHUNTER), maps of the region were
constructed based on Krauter *et al.*, 1995 (Krauter, K., *et al.*, *Nature* 377:321-333 (1995)). The
30 map used is shown in Figure 1, which indicates the markers used in the present study and those
reported in Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241
(1997)).

In the interest of clarity and simplicity, nominal p-values are reported here. Most of the
methods used involve multiple comparisons (*e.g.*, testing across multiple genetic models, or using

multiple different approaches to the same question). Therefore, only p-values of 0.01 or less should be viewed as significant (Pocock, S., *et al.*, *N. Engl. J. Med.* 317:426-432 (1987)).

Genetic linkage analysis in the sample of 286 families was performed next for *A2M* and a set of chromosome 12 markers (Figure 1) including several centromeric markers recently reported to be linked to AD by Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)). *A2M* resides roughly 27 cM of the centromeric marker, D12S1042, which yielded the highest maximum 2-point lod-score (2.7) by affecteds-only analysis in the Pericak-Vance *et al.* study ((Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997); Krauter, K., *et al.*, *Nature* 377:321-333 (1995)). Besides setting out to determine whether the association between *A2M-2* and AD could be detected by traditional linkage analyses and could be explained by the previous linkage findings of Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)), the inventors also attempted to confirm the previous linkage findings in the sample set.

Initially affecteds-only lod score analyses were performed, which showed that lod scores for *A2M* and virtually all markers were less than 0 out to a recombination fraction of 30 cM. Lod scores from affecteds only analyses were also strongly negative in Tier 3 (in which at least one affected in each family does not carry the *APOE-ε4* allele). Tier 3 is analogous to the subgroup of AD families for which Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)) reported evidence for linkage to AD. The lod scores were similarly negative in Tier 2 (families in which all affecteds are *APOE-ε4* carriers, but at least one is a non-homozygote). However, in Tier 1 (families in which all affecteds are *APOE-ε4/ε4*), there was a weak but non-significant signal: maximum lod score of 1.23 at $\Theta = 7$ cM for D12s1042 and .74 at $\Theta = 12$ cM for D12s1090, 7cM distal. The results of several different age curve models (not shown), including that used by Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)), were also strongly negative, except in Tier 1, where the results were slightly positive yet nonsignificant as in the affecteds-only analyses.

Next, multipoint analyses were performed using GENEHUNTER (Kruglyak, L., *et al.*, *Am J Hum Genet* 58:1347-1363 (1996))(software available from Kruglyak *et al.* by anonymous ftp at ftp-genome.wi.mit.edu or from their World Wide Web site <http://www-genome.wi.mit.edu/ftp/distribution/software/genehunter>)) (Figures 2A-2D) and the results were

5 similarly negative. NPL scores for *A2M* and virtually all markers were less than 0 for the sample as a whole (Figure 2A) and for Tiers 2 (Figure 2C) and 3 (Figure 2D) (analogous to the subgroup analyzed by Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)). However, in Tier 1 (Figure 2B), in which all affecteds are *APOE-ε4/ε4*, there was again a weak but non-significant signal: NPL = 1.37 at D12sl042 ($p = .08$) and 1.21 at D12sl090 ($p = A1$).

10 Multipoint sibpair analyses in the program ASPEX were similarly negative with maximum lod scores of 1.31 for D12sl042, and .91 for D12sl090. Of note, exclusion analyses with "risk" (corresponding approximately to λ , the recurrence risk in relatives) set at 2 performed in ASPEX yielded extremely negative lod scores throughout the region for the sample as a whole ($\text{lod} < -8$), and for Tiers 2 and 3 ($\text{lod} < -3$). For Tier 1, consistent with the non-significant peak observed in several analyses, the region provided no significant results for genetic linkage, but could not be entirely excluded.

15 Next, all of the above analyses were repeated using uncorrected allele frequencies. The analyses defining the sample was also repeated as in Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)), *i.e.*, using only the 259 families in which all sampled affected individuals experienced onset of AD at 60 or above, and in which at least 3 cases of AD spanning 2 generations were evident. The results of all of these analyses were similar to those described above, and did not confirm the findings presented in Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)). Thus, similarly negative findings were obtained using multiple genetic models, multiple family selection criteria, and multiple statistical methods. In all cases, the inventors were unable to confirm the previous finding of linkage of AD to the centromeric region of chromosome 12 reported by Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)). It should also be noted that traditional linkage and multipoint analyses were not able to detect genetic linkage between *A2M* and AD. This is most likely due to the lack of informativeness of the biallelic *A2M* polymorphism in a sample of this nature. However, evidence was found for linkage of chromosome 12 to AD using other markers, including *D12598* and *DRS358*, (Wu, W. S., *et al.*, *JAMA* (in press, 1998)) that were located much closer to *A2M* than the markers used in the genetic analysis described above.

5 C. Discussion

10 These analyses provide evidence of a link between *A2M* and risk for Alzheimer's disease, and demonstrate that inheritance of the *A2M-2* allele, confers three- to four-fold increased risk for AD and is strongly associated with AD in the SDT. Moreover, in these analyses, the inheritance of one or two *A2M-2* alleles; conferred a degree of risk for AD comparable to that associated with inheriting two *APOE-ε4* alleles. The association of *A2M-2* and AD was not accounted for by differences in the dose of the *APOE-ε4* allele, and did not vary when the *APOE-ε4* effect was controlled for in a multivariate setting.

15 Although, *A2M* maps within 30 cM of the chromosome 12 markers reported to be linked to AD (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997); Krauter K., *et al.*, *Nature* 377:321-333 (1995)), the observed genetic association between *A2M* and AD described here does not appear to account for these linkage findings. Moreover, the inventors were unable to confirm the linkage reported by Pericak-Vance *et al.* (Pericak-Vance, M. A., *et al.*, *JAMA* 278(15):1237-1241 (1997)) in the present data set despite rigorous attempts to duplicate that study. The inventors were also unable, using traditional genetic linkage and multipoint analyses, to demonstrate genetic linkage of *A2M* to AD, which may reflect the limited power of such analyses in a data set of this nature, especially in comparison to family-based association methods in the analysis of a candidate gene.

25 Although these data suggest that *A2M-2*'s association with AD may be as strong as that of *APOE-ε4*, *A2M-2* does not appear to share *APOE-ε4*'s effect on age of onset of AD (based on the similarity in age of onset among affecteds with different *A2M* genotypes seen graphically and analytically). Thus, rather than conferring risk by modifying the age of onset, as appears to be the case with *APOE-ε4*, it appears that *A2M-2* confers risk for developing the disease at any age. The mode of inheritance for the observed effect of *A2M-2* on AD could not be assessed in the present study. For example, it is unclear whether *A2M* acts alone or in conjunction with other genes to confer increased risk for AD. In addition, due to the small number of *A2M-2* homozygotes, the inventors were unable to assess for a difference in effect of one versus two doses of *A2M-2* on risk for AD.

5 Since increased *APOE-ε4* dose correlated with an earlier age of onset among AD patients with the *A2M-2* allele, it is possible that *APOE-ε4* influences the age of onset in individuals who are a priori more susceptible to developing AD due to the inheritance of *A2M-2*. *A2M-2* may represent a relatively prevalent public polymorphism (25-30%) that confers susceptibility for AD, with age of onset of the disease in susceptible individuals modified by *APOE-ε4* dosage. Further investigation will be necessary to identify other genes that may epistatically interact with *A2M* and/or *APOE* to determine either risk and/or the age of onset of AD. The genetic model suggested by these current findings is in agreement with the recent observations of Meyer *et al.* (Submitted), in a study of nearly 5,000 elderly individuals, that APOE genotype influences "when" but not "whether" AD will develop in a priori susceptible individuals, and that regardless of APOE genotype, roughly 50% of the population will not develop AD by age 100. In view of these findings, inheritance of mutations in genes like *A2M* or other genes yet to be identified may be necessary to confer initial predisposition for the occurrence of AD, while genes like *APOE* modify age of onset in susceptible individuals.

20 This heuristic genetic model for AD lends itself to an intriguing biological model for AD neuropathogenesis in view of the known physiological roles of α_2 M, apoE, APP, and their common receptor, LRP. α_2 M is a pan-protease inhibitor which is constitutively expressed in brain and upregulated together with *LRP* during acute phase brain injury (e.g., by ischemia, stroke). α_2 M has been shown to bind a wide range of proteases, growth factors, cytokines, and small polypeptides, and has been detected in pyramidal neurons and neuritic plaques in affected regions of AD brain (Borth, W., *FASEB* 6: 3345-3353 (1992); Bauer J., *et al.*, *FEBS* 285:111-114 (1991)). Recently, α_2 M was shown to tightly bind A β and mediate its clearance via endocytosis through LRP (Narita, M., *et al.*, *J. Neurochem.* 69:1904-1911 (1997)), and, in complex with a serine protease, has been shown to degrade A β (Qiu, W. Q., *et al.*, *J. Biol. Chem.* 271:8443-8451 (1996)). Thus, α_2 M may clear and/or degrade secreted A β from the brain parenchyma, thereby governing the rate of A β fibrillogenesis and β -amyloid deposition. Indeed, α_2 M has also been shown to attenuate both A β fibril formation and neurotoxicity *in vitro* (Du, Y., *et al.*, *J. Neurochem.* 69:299-305 (1997); Hughes, S. R., *et al.*, *Proc. Natl. Acad. Sci. (USA)* 95:3275-3280 (1998); Du, Y., *et al.*, *J. Neurochem.* 70:1182-1188 (1998)). Since apoE, like α_2 M, can bind A β and utilizes LRP as its major neuronal receptor in brain, these two proteins could

5 conceivably compete for binding to LRP and A β . Thus, the clearance and degradation of A β in the brain parenchyma could potentially be impaired by increased amounts of apoE (Zhang, Z., *et al.*, *Int. J. Exp. Clin. Invest.* 3:156-161 (1996)), or the presence of apoE isoforms that have higher affinity for LRP or A β .

10 Interestingly, *APOE* promoter polymorphisms, which upregulate transcription of *APOE*, have recently been shown to be associated with AD and to be in linkage disequilibrium with *APOE- ϵ 4* (Bullido, M. J., *et al.*, *Nature Genet.* 18:69-71 (1998); Lambert, J.C., *et al.*, *Hum. Mol. Genet.* 7:533-540 (1998)). Additionally, allelic distortion resulting in higher levels of message for *APOE- ϵ 4* (relative to *APOE-3*) has been reported to occur in the brains of AD patients but not age-matched controls (Lambert, J., *et al.*, *Hum. Mol. Genet.* 6:2151-2154 (1997)). The absence of apoE in transgenic mice expressing an FAD mutant form of *APP* has been shown to attenuate β -amyloid deposition (Bales, K.R., *et al.*, *Nat. Genet.* 17:263-264 (1997)). Given these findings, it is possible that increased levels of *APOE* expression may modify age of onset of A β perhaps by interfering with α_2 M-mediated clearance and degradation of AD via LRP-mediated endocytosis and direct degradation by α_2 M-protease complexes, respectively, resulting in greater amyloid burden. The inventors have recently localized LRP to synaptic terminals where impaired clearance of α_2 M complexes (*e.g.*, containing A β) could conceivably promote synaptic degradation, a major neuropathological feature of AD (Kim *et al.*, Submitted). A heuristic genetic model in which *A2M-2* confers susceptibility to AD while *APOE* genotype modifies the age of onset readily lends itself to this biological model. In further support for this model, an exon 3 polymorphism in *LRP* has also recently been shown and confirmed to be genetically associated with AD (Kang, D.E., *et al.*, *Neurology* 49:56-61 (1997); Lendon, C. L., *et al.*, *Neurosci. Lett.* 222:187-190 (1997); Wavrant-DeVrieze, F., *et al.*, *Neurosci. Lett.* 227:68-70 (1997); Hollenbach, E., *et al.*, *Neurology*, In Press (1998)). However, another polymorphism consisting of a trinucleotide repeat in the 5' untranslated region of *LRP* was excluded from being genetically linked or associated with AD (Lendon, C.L., *et al.*, *Neurosci. Lett.* 222:187-190 (1997); Clatworthy, A.E., *et al.*, *Arch. Neurol.* 54:1289-1292 (1997); and Scott, W.K., *et al.*, *Neurogenetics* 1:179-183 (1998)).

30 The data above indicate that *A2M* is a novel AD gene that predisposes, in particular, carriers of the *A2M-2* deletion to increased risk for AD, but without modifying age-of-onset.

5 *A2M*, together with the gene for its receptor, *LRP*, and the genes encoding two other LRP ligands, *APOE*, and *APP*, have now all been genetically linked to AD. Thus it is plausible that all four of these proteins participate in a common pathogenic pathway leading to AD-related neurodegeneration.

10 The *A2M-2* exon splice site deletion occurs at the 5' splice acceptor for exon 18. One would predict that this deletion would cause the skipping or deletion of exon 18, which encodes the second half of the bait domain of $\alpha 2M$. The bait domain is required for $\alpha 2M$ to bind and trap a wide range of proteases. The inventors postulate that the loss of a fully functional bait domain in a portion of $\alpha 2M$ molecules would result in the promotion of AD-related neuropathogenesis in a variety of different ways. One possibility is that the loss of the
15 bait domain results in decreased ability to degrade and clear $A\beta$ from the brain (*e.g.*, in synapses). However, the possible consequences of the loss of a fully functional bait domain are actually immense in scope given the incredibly wide range of proteases that $\alpha 2M$ can bind, and the multifold downstream events which may occur as a result of the attenuation of global protease regulation (*e.g.*, following injury to the brain due to stroke or head trauma). The location of the
20 *A2M-2* exon splice site deletion suggests that the protease inhibitory function of $\alpha 2M$ may protect against AD. Thus, the possibility that any of the multifold functions of $\alpha 2M$ may be adversely affected by *A2M-2* cannot be ruled out. For example, in addition to binding to LRP, $\alpha 2M$ can also bind to the $\alpha 2M$ signalling receptor ($\alpha 2MSR$), which is involved in cell signalling events (*e.g.*, activation of phosphatidylinositol 3-kinase) modulating intracellular calcium levels (Misra, U. K.
25 and Pizzo, V., *J. Biol. Chem.* 273:13399-13402 (1998)).

Example 2

The G Allele

A. Genotype and Allele Frequencies

An initial exploratory experiment was used to test for the possibility that the less common "G" allele of *A2M* would be associated with Alzheimer's disease. Because of the potential for error due to multiple hypothesis testing, an initial data set was used as an exploratory data set, with the intent of formulating specific hypotheses that would then be tested formally with a second independent data set.

In the initial data set, 90 non-Alzheimer's individuals, age-compatible, who had either undergone screening tests with the Blessed dementia scale (Blessed, G., *et al.*, *Br. J. Psychiatry* 114:797-811 (1968)) or whose DNA had been isolated from autopsy material and were demonstrated to not have Alzheimer's disease, and 171 individuals who either had a clinical diagnosis of Alzheimer's disease or neuropathologically proven Alzheimer's disease were genotyped (Table 6). The 90 control subjects were primarily spouses of the AD subjects.

Genotypes were determined by polymerase chain reaction amplification of DNA and RFLP analysis using MboI (Figure 3). Genomic DNA isolated from brain tissue and blood was amplified by polymerase chain reaction (PCR) in the presence of oligonucleotide sense primer C23 (SEQ ID NO: 3) and antisense primer AS24 (SEQ ID NO: 4), 10 mM Tris-HCl, 50 mM KCl (pH 8.3), 1.5 mM MgCl₂, 5 mM dNTPs, 5 pmol each primer, and 1.25 U *Taq* DNA polymerase. The PCR was carried out in a touchdown procedure that stepped down the annealing temperature to increase primer specificity as follows: 1 cycle at 94°C-5 min; 4 cycles at 94°C-30 sec, 65°C-30 sec, 72°C-1 min; 4 cycles at 94°C-30 sec, 62°C-30 sec, 72°C-1 min; 4 cycles at 94°C-30 sec, 59°C-30 sec, 72°C-1 min; 20 cycles at 94°C-30 sec, 56°C-30 sec, 72°C-1 min; and 1 cycle at 72°C-5 min. For each reaction mixture, 10 units of MboI were added to the amplified product of 615 bps and digestion carried out at 37°C for 3 hours, producing fragments of 532 and 429 bps. The digested product was loaded onto a 2% agarose gel treated with ethidium bromide (0.005%) and electrophoresed for 2 hours under

5 constant voltage (150V), which is sufficient to separate the digested product so that the 532 and 429 bp bands can be distinguished. After electrophoresis, DNA fragments were visualized by UV illumination using a Biorad Geldoc system. Incomplete digestion was monitored by looking for the presence of the 615 bp fragment.

10 For the statistical analyses below, comparisons of *A2M* allele frequency (proportion of chromosomes in which an allele is present) and genotype frequency (proportion of individuals with a genotype) were performed with 2x2 tables using Fisher's exact test for significance. Age of onset of Alzheimer's disease was normally distributed and compared by t-test. Multivariate analysis for odds of AD was performed by logistic regression with *APOE* genotype coded according to the presence or absence of *APOE-ε4*. Similar results were
15 obtained when the group with *APOE-ε4* was coded separately as heterozygotes or homozygotes for this allele. Odds ratios are presented with 95% confidence intervals (CI). All analyses were performed with Stata software (Stata Corporation, College Park, Texas). All significance tests were two-tailed.

20 The results of the genotype and allele frequency analysis for the exploratory data set suggested that the *A2M* Val/Ile¹⁰⁰⁰ polymorphism may be a genetic risk factor for Alzheimer's disease, and warranted further investigation. In the control series the G allele frequency was 0.28 and 6 of 90 individuals contained the *A2M* G/G genotype (0.067). In the Alzheimer's disease set, the G allele frequency was 0.32. The *A2M* G/G genotype frequency was increased
25 in the Alzheimer's group at 0.12 (Table 6).

Table 6: Overrepresentation of the *A2M* G/G genotype in AD

Exploratory Data Set	N	G allele (%)	G/G genotype (%)
Control	90	0.28	0.07
AD	171	0.32	0.12
Hypothesis Testing Data set			
Control	359	0.32	0.07
AD	566	0.34	0.12*
Combined Data Set (Total)			
Control	449	0.32	0.07
AD	737	0.34	0.12**

* $p < 0.05$

** $p < 0.01$

To formally test the hypothesis that the *A2M* G/G genotype of *A2M* is overrepresented in Alzheimer's disease, another data set was collected and genotyped from additional independent groups of patients and controls. Power analysis showed that more than 500 AD and control individuals would be necessary to have an 80% chance of showing a difference between genotype frequencies of 0.07 and 0.12. Therefore, AD patients and control individuals collected from several sites were used to form a sample of this size. This second data set consisted of individuals who met the criteria for AD or control individuals used for the initial data set, and included 387 sporadic AD and 359 control individuals collected from the Massachusetts General Hospital Memory Disorder Unit and from a consortium of European Centers (University of Hamburg, Germany; University of Basel, Switzerland; University of Brescia, Italy) and 179 probands from NIMH Genetics Initiative sample (Blacker, D., *et al.*, *Neurology* 48:139-47 (1997)).

The results from the second data set supported the hypothesis, with G/G genotype 0.07 in controls and 0.12 in AD ($p=0.05$, Fisher's exact test) (Table 6). The G allele was not over-represented in AD in this data set (control = 0.32, AD = 0.34) (Table 6). No difference in age of onset between G/G and non G/G was found in the second data set. Multivariate analysis showed that site of collection did not influence genotype frequencies. In addition,

5 when the rate in controls was compared to the subset of 387 clinic based AD cases (G/G frequency = 11.4%, $p=0.06$) and the subset of 179 familial AD samples (G/G frequency = 12.7%, $p=0.056$), essentially equal overrepresentation of G/G was found in each. These data support an association of the G/G genotype with AD.

10 Next, a series of exploratory analyses were done using pooled data from both data sets (737 AD patients and 449 controls). The G/G genotype was present in 11.9% of the Alzheimer patients, and 7.3% of the controls ($p<0.01$, Fisher's exact test). The genotype frequency in controls was consistent with Hardy Weinberg equilibrium. Age of onset was not different between AD patients with G/G (70.0 ± 9.2 , $X\pm SD$) and non-G/G carriers (70.6 ± 9.1).

15 **B. Multivariate Model Controlling for Presence of APOE- $\epsilon 4$ and Gender**

A multivariate logistic model that controlled for the presence of the APOE- $\epsilon 4$ allele and gender showed an odds ratio of 1.88 (95% CI 1.20-2.95; $p<0.01$) for the presence of A2M G/G genotype. The presence of APOE- $\epsilon 4$ in this model was associated with an odds
20 ratio of 4.21 (95% CI 3.20-5.55; $p<0.001$). The odds ratio for the combination of A2M G/G and APOE- $\epsilon 4$ is 9.68 (95% CI 3.91-24.0; $p<0.001$) relative to those with neither risk factor.

To explore whether the A2M G/G mediated risk was influenced by the presence or absence of APOE- $\epsilon 4$, the sample was stratified by the presence or absence of APOE- $\epsilon 4$. Similar overrepresentations of the A2M G/G genotype were seen in the strata with or without
25 APOE- $\epsilon 4$, suggesting that the effects of the two risk factors are independent. Multivariate analysis for interaction between the A2M G/G genotype and either APOE- $\epsilon 4$ or gender demonstrated no interactions ($p>0.5$ for interaction terms). Analysis based on the Mantel-Haenzel estimator suggested no heterogeneity of odds ratios between strata with and without APOE- $\epsilon 4$ ($p>0.5$), consistent with the absence of an interaction between A2M and APOE- $\epsilon 4$
30 in our logistic regression model.

5 C. *Biological and Neuropathological Effects of G/G Genotype*

Also investigated were the possible biological effects of inheritance of the *A2M* G/G genotype. Previous studies of *A2M* immunohistochemistry performed on Alzheimer's brain suggested that senile plaques in different patients stained with varying intensity (Rebeck, G. W., *et al.*, *Ann. Neurol.* 37:211-217 (1995)). The possibility that this variability was due to different genotypes was investigated by immunostaining samples from 8 AD cases known to be G/G or A/A genotype.

10 Sections of brain tissue from AD patients were incubated with anti- α_2 M antibody (1:500) from Zymed. A goat Cy-3 linked secondary antibody (Jackson Immunoresearch) was used to visualize immunostaining. In some instances, double immunofluorescence was carried out with an A β counterstain (antibody 10D5, obtained from Dr. D. Schenk, Athena Neurosciences) using bodipyfluorescein linked secondary antibody (Molecular Probes) as the second fluorochrome. In all instances, α_2 M immunoreactivity was present robustly on senile plaques, astrocytes, and neurons, and no qualitative differences were observed between the genotype groups.

20 Next explored was the effect inheritance of the *A2M* G/G genotype on the neuropathological phenotype of Alzheimer's disease. In an earlier study, reported in Gomez-Isla *et al.* (Gomez-Isla, T., *et al.*, *Ann. Neurol.* 41:17-24 (1997)), stereological techniques and quantitative image analysis were used to measure the amount of amyloid burden present in neocortical association area surrounding the superior temporal sulcus (STS) and the number of neurofibrillary tangles present in the same region in normal and Alzheimer brain. Thirty-one (31) of the previously analyzed AD cases (Gomez-Isla, T., *et al.*, *Ann. Neurol.* 41:17-24 (1997)) were genotyped and selected for further analysis. Initial inspection of the data suggested an increase in A β in individuals who were A/G or G/G, with no difference between these two groups. A statistically significant increase in A β deposition was seen comparing the G containing cases to the non-G containing cases ($8.8 \pm 2.8\%$ vs $6.6 \pm 1.9\%$, $p < 0.03$, unpaired t test) (Table 7). The effect appears to be primarily due to the presence of at least one G allele, in that no difference between A/G and G/G genotypes was observed although the sample size limits the power of this comparison. There were no statistically

- 5 significant differences in neurofibrillary tangle number with G alleles, suggesting a specific effect on A β deposition.

Table 7: Neuropathological Correlates of *A2M* Genotype

	Genotype	n	Amyloid burden (%)	Neurofibrillary Tangles(x10 ³)
10	A/A	10	6.6 \pm 1.9*	6.9 \pm 4.1
	A/G + G/G	21	8.8 \pm 2.8**	9.0 \pm 5.0***

* mean \pm SD

** p<0.03 t test

15 *** n=17 for this measurement

D. Discussion

- 20 These analyses provide further support for an association between *A2M*, and in particular the *A2M* G/G genotype, and AD. In order to overcome some of the difficulties inherent in exploratory studies of genetic risk factors, independent data sets were used to first generate, then test the hypothesis that the G/G genotype is associated with Alzheimer's disease. Exploratory studies of genetic risk factors run the risk of Type I errors because of the
- 25 large number of hypotheses that are either explicitly or implicitly being tested. Alternatively, type II errors can occur if sample size is insufficient. The two separate control populations, used to overcome these difficulties, gave identical genotype frequencies. Moreover, the hypothesis testing data set was itself made up of two groups (sporadic AD and probands from a multicenter study of Alzheimer's disease sib pairs and small families) each of which gave
- 30 essentially identical genotype frequencies and overrepresentation of the G/G genotype. The *A2M* G/G effect was not dependent on *APOE* genotype, and the effect was additive and substantial -- the odds ratio of G/G in the presence of an *APOE*- $\epsilon 4$ allele showed an almost 10 fold increased risk. Power analysis using the size of effect observed in our combined data set (using about 500 samples) suggests that a group size of 578 controls and 578 Alzheimer's

5 disease patients would be necessary to have an α of 0.05 and a β of 80% for the hypothesis that the G/G genotype is increased from 0.07 to 0.12 in Alzheimer's disease.

The association of the A2M G/G genotype as a risk factor in Alzheimer's disease leads to several interesting mechanistic interpretations. Previous studies have shown that α_2 M immunostains SP and have implicated α_2 M in amyloid precursor protein metabolism (Ganter, U., *et al.*, *FEBS* 282:127-131 (1991)) as well as possible interactions with binding A β (Du, Y., *et al.*, *J. Neurochem.* 69:299-305 (1997)) and mediating degradation of A β (Qiu, W. Q., *J. Biol. Chem.* 271:8443-8451 (1996); Zhang, Z., *et al.*, *Int. J. Exp. Clin. Invest.* 3:156-161 (1996)). α_2 M-A β complexes undergo endocytosis via α_2 M-r/LRP (Narita, M., *et al.*, *J. Neurochem.* 69:1904-1911 (1997)). α_2 M associates with A β and prevents fibril formation (Hughes, S., *et al.*, *Proc. Natl. Acad. Sci.* 95:3275-3280 (1998)). In addition, α_2 M can be viewed as a (sub) acute phase reactive molecule and as a potent protease inhibitor which might participate in the inflammatory response in Alzheimer brain. Interestingly, α_2 M has also been implicated as a trophic factor, possibly as a result of binding and transporting a number of polypeptide growth factors and cytokines (Borth, W., *FASEB* 6:3345-3353 (1992)). Like Apo E, α_2 M may serve as a binding protein for targeting various bioactive molecules to their site of action or clearing them (Rebeck, G.W., *et al.*, *Ann. Neurol.* 37:211-7 (1995)). Moreover, the fact that apoE and APP, two other LRP ligands are clearly involved in the pathophysiology of Alzheimer's disease, supports the hypothesis that a link exists between A2M and risk for Alzheimer's disease. In fact, a polymorphism in LRP has recently been reported in two studies showing a genetic association with Alzheimer's disease (Kang, D. E., *et al.*, *Neurology* 49:56-61 (1997); Hollenbach, E., *et al.*, *Neurology* 50:1905-7 (1998)).

The likelihood that A2M is a risk factor is also supported by the association between the A2M-2 allele and AD discussed in Example 2 above. The combined analyses of the A2M-2 and the A2M-G alleles indicate that these two mutations act independently of each other to confer an increased risk of AD, *i.e.*, inheritance of A2M-2 is not genetically linked to inheritance of A2M-G in the same AD patients. This is based on the observation that the A2M-G allele frequently segregates with the A2M-1 allele. In 160 random individuals, 56 of 58 obligate A2M-G chromosomes were found to carry the A2M-1 allele (13 out of 14 obligate A2M-2 chromosomes carry the A2M-A allele).

5 In summary, *A2M* appears to be a novel AD gene that predisposes, in particular, individuals with either the *A2M*-2 or the *A2M* G/G genotype to increased risk for AD, but without modifying age-of-onset.

10 Modifications of the above-described modes for carrying out the invention that are obvious to persons of skill in medicine, genetics, molecular biology, biochemistry and/or related fields are intended to be within the scope of the following claims.

 All publications and patents mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patents are herein incorporated by reference to the same extent as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

15 Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications can be practiced within the scope of the appended claims.

5 ***What Is Claimed Is:***

1. A method of diagnosing Alzheimer's disease comprising the step of genotyping the Alpha-2-Macroglobulin locus of an individual.

10 2. The method of claim 1, wherein said genotyping step comprises the steps of:
 (a) isolating nucleic acid from an individual;
 (b) amplifying the nucleic acid to generate an *A2M* fragment; and
 (c) analyzing the fragment thereby correlating *A2M* genotype with the
occurrence of Alzheimer's disease.

15 3. The method of claim 2, wherein the nucleic acid is DNA.

4. The method of claim 2, wherein the nucleic acid is RNA.

20 5. The method of claim 2, wherein said step (b) utilizes polynucleotide primers X and Y, wherein said primers X and Y anneal to nucleotides flanking the site of the mutation present in the *A2M-2* allele.

25 6. The method of claim 2, wherein said step (b) utilizes polynucleotide primers X and Y, wherein said primers X and Y anneal to nucleotides flanking the site of the mutation present in the *A2M-G* allele.

7. The method of claim 2, wherein said step (c) comprises sequencing the fragment to determine *A2M* genotype.

30 8. The method of claim 2, wherein said step (c) comprises RFLP analysis of the fragment to determine *A2M* genotype.

5 9. The method of claim 2, wherein said step (c) comprises size fractionation of the fragment to determine *A2M* genotype.

 10. The method of claim 5, wherein said step (c) comprises sequencing the fragment to determine *A2M* genotype.

10 11. The method of claim 5, wherein said step (c) comprises RFLP analysis of the fragment to determine *A2M* genotype.

 12. The method of claim 5, wherein said step (c) comprises SSCP analysis of the fragment to determine *A2M* genotype.

15 13. The method of claim 6, wherein said step (c) comprises sequencing the fragment to determine *A2M* genotype.

20 14. The method of claim 6, wherein said step (c) comprises RFLP analysis of the fragment to determine *A2M* genotype.

 15. The method of claim 6, wherein said step (c) comprises SSCP analysis of the fragment to determine *A2M* genotype.

25 16. The method of claim 1, wherein said genotyping step comprises the steps of:
 (a) isolating DNA from an individual
 (b) subjecting said DNA to RFLP analysis thereby correlating *A2M*
genotype with the occurrence of Alzheimer's disease.

30 17. The method of claim 16, wherein said RFLP analysis utilizes a restriction endonuclease specific for a restriction site created or deleted due to the pentanucleotide deletion found in *A2M-2*.

5 18. The method of claim 16, wherein said RFLP analysis utilizes a restriction endonuclease specific for a restriction site created or deleted due to the substitution mutation found in *A2M-G*.

10 19. A method for diagnosing Alzheimer's disease comprising: isotyping the Alpha-2-Macroglobulin protein of an individual.

 20. The method of claim 19 comprising the steps of:
 (a) isolating protein from said individual
 (b) analyzing the protein thereby correlating Alpha-2-Macroglobulin
15 isotype with the occurrence of Alzheimer's disease.

 21. The method of claim 20, wherein said step (b) comprises western blot analysis of the protein to determine *A2M* genotype.

20 22. The method of claim 20, wherein said step (b) comprises ELISA analysis of the protein to determine *A2M* genotype.

 23. The method of claim 20, wherein said step (b) comprises α_2 M electrophoretic mobility assay analysis of the protein to determine *A2M* genotype.

25 24. The method of claim 21, wherein said western blot analysis utilizes an antibody specific for the α_2 M-2 variant.

 25. The method of claim 21, wherein said western blot analysis utilizes an antibody
30 specific for the α_2 M Val¹⁰⁰⁰ variant.

 26. The method of claim 22, wherein said ELISA analysis utilizes an antibody specific for the α_2 M-2 variant.

5 27. The method of claim 22 wherein said ELISA analysis utilizes an antibody specific for the α_2 M Val¹⁰⁰⁰ variant.

10 28. A diagnostic kit for diagnosing Alzheimer's disease according to the method of claim 2, comprising: (i) a carrier means compartmentalized in close confinement therein to receive one or more container means; (ii) a container means containing polynucleotide primers X and Y, wherein said primers X and Y anneal to nucleotides flanking the site of the mutation present in the *A2M-2* allele; (iii) a container means containing α_2 M-1 DNA, or fragment thereof; and (iv) a container means containing α_2 M-2 DNA, or a fragment thereof.

15 29. A diagnostic kit for diagnosing Alzheimer's disease according to the method of claim 2, comprising: (i) a carrier means compartmentalized in close confinement therein to receive one or more container means; (ii) a container means containing the polynucleotide primers X and Y, wherein said primers X and Y anneal to nucleotides flanking the site of the mutation present in the *A2M-G* allele; (iii) a container means containing α_2 M Ile¹⁰⁰⁰ DNA, or fragment thereof; and (iv) a container means containing α_2 M Val¹⁰⁰⁰ mutant DNA, or a fragment thereof.

25 30. A diagnostic kit for diagnosing Alzheimer's disease according to the method of claim 20, comprising: (i) a carrier means compartmentalized in close confinement therein to receive one or more container means; (ii) a container means containing an antibody specific for the α_2 M-2 variant; (iii) a container means containing an antibody specific for α_2 M-1; (iv) a container means containing the α_2 M-2 variant, or fragment thereof; and (v) a container means containing α_2 M-1, or fragment thereof.

30 31. A diagnostic kit for diagnosing Alzheimer's disease according to the method of claim 20, comprising: (i) a carrier means compartmentalized in close confinement therein to receive one or more container means; (ii) a container means containing an antibody specific for the α_2 M Val¹⁰⁰⁰ variant; (iii) a container means containing an antibody specific for the α_2 M

- 5 Ile¹⁰⁰⁰ protein; (iv) a container means containing the α_2 M Val¹⁰⁰⁰ variant, or fragment thereof; and (v) a container means containing the α_2 M Ile¹⁰⁰⁰ protein, or fragment thereof.

- 10 32. A diagnostic kit for diagnosing Alzheimer's disease according to the method of claim 23, comprising: (i) a carrier means compartmentalized in close confinement therein to receive one or more container means; (ii) a container means containing a protease; (iii) a container means containing a substantially purified sample of the α_2 M-2 variant; and (iv) a container means containing a substantially purified sample of the fast form of α_2 M-1.




FIG.1

SUBSTITUTE SHEET (RULE 26)




FIG. 2A




FIG. 2B

SUBSTITUTE SHEET (RULE 26)




FIG. 2C




FIG. 2D

SUBSTITUTE SHEET (RULE 26)

4/4




FIG.3

-1-

SEQUENCE LISTING

<110> Tanzi, Rudolph E.
Hyman, Bradley T.
Rebeck, George W.

<120> An Alpha-2 Macroglobulin Diagnostic Test

<130> 0609.446PC02

<140> To be assigned

<141> 1998-09-04

<150> US 60/093,297

<151> 1998-07-17

<150> US 60/057,655

<151> 1997-09-05

<160> 4

<170> PatentIn Ver. 2.0

<210> 1

<211> 23

<212> DNA

<213> Homo sapiens

<400> 1

ctttccttga tgaccaagc gcc

23

<210> 2

<211> 22

<212> DNA

<213> Homo sapiens

<400> 2

gttgaaaata gtcagcgacc tc

22

<210> 3

<211> 20

<212> DNA

<213> Homo sapiens

<400> 3

atccctgaaa ctgccaccaa

20

<210> 4

<211> 20

<212> DNA

<213> Homo sapiens

<400> 4

gtaactgaaa cctactggaa

20

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/18535

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 7.1, 91.1, 91.2; 536/23.1, 24.3, 24.31, 24.33; 530/350, 387.1, 388.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	BLACKER et al. Alpha-2 macroglobulin is genetically associated with Alzheimer disease. Nature Genetics. 19 August 1998, Vol. 19, pages 357-360, see entire document.	1-32
A, P	PERICAK-VANCE et al. Complete genomic screen in late-on-set familial Alzheimer disease. JAMA. 15 October 1997, Vol. 278, No. 15, pages 1237-1241, see entire document.	1-32
A, P	BLACKER et al. The Genetics of Alzheimer disease. Arch. Neurol. March 1998, Vol. 55, pages 294-296, see entire document.	1-32
A	SHELLENBERG. G.D. Genetic dissection of Alzheimer disease, a heterogeneous disorder. Proc. Natl. Acad. Sci. September 1995, Vol. 92, pages 8552-8559, see entire document.	1-32

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 NOVEMBER 1998

Date of mailing of the international search report

23 DEC 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JEFFREY FREDMAN

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet) (July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/18535-

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	POLLER et al. Sequence polymorphism in the human alpha-2-macroglobulin (A2M) gene. Nucleic Acids Research. 1991, Vol. 19, page 198, see entire document.	28, 29
X	JUSTUS et al. Quantification of free alpha 2-macroglobulin and alpha 2-macroglobulin protease complexes by a novel ELISA system based on streptococcal alpha 2 macroglobulin receptors. Journal of Immunological Methods. 24 January 1990, Vol. 126, pages 103-108, see entire document.	30-32

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/18535

A. CLASSIFICATION OF SUBJECT MATTER: IPC (6):

C12Q 1/68; C12P 19/34; G01N 33/54; C07H 21/02, 21/04; C07K 15/26, 15/28

A. CLASSIFICATION OF SUBJECT MATTER: US CL :

435/6, 7.1, 91.1, 91.2; 536/23.1, 24.3, 24.31, 24.33; 530/350, 387.1, 388.1

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, CAPLUS, MEDLINE, BIOSIS, BIOTECHABS, WPIDS

search terms: alpha, macroglobulin, alzheimer, PCR, polymerase, chain, antibody, eliza, allele, mutation, mutant, variant, alteration, monoclonal, antigen

